

# Black hole size revealed by its eating pattern

August 12 2021

---



An artist's impression of an accretion disk rotating around an unseen supermassive black hole. The accretion process produces random fluctuations in luminosity from the disk over time, a pattern found to be related to the mass of the black hole in a new study led by University of Illinois Urbana-Champaign researchers. Credit: Mark A. Garlick/Simons Foundation

The feeding patterns of black holes offer insight into their size,

researchers report. A new study revealed that the flickering in the brightness observed in actively feeding supermassive black holes is related to their mass.


Supermassive [black holes](#) are millions to billions of times more massive than the sun and usually reside at the center of massive galaxies. When dormant and not feeding on the gas and stars surrounding them, SMBHs emit very little light; the only way astronomers can detect them is through their gravitational influences on stars and gas in their vicinity. However, in the [early universe](#), when SMBHs were rapidly growing, they were actively feeding—or accreting—materials at intensive rates and emitting an enormous amount of radiation—sometimes outshining the entire galaxy in which they reside, the researchers said.

The new study, led by the University of Illinois Urbana-Champaign astronomy graduate student Colin Burke and professor Yue Shen, uncovered a definitive relationship between the mass of actively feeding SMBHs and the characteristic timescale in the light-flickering pattern. The findings are published in the journal *Science*.

The observed light from an accreting SMBH is not constant. Due to [physical processes](#) that are not yet understood, it displays a ubiquitous flickering over timescales ranging from hours to decades. "There have been many studies that explored possible relations of the observed flickering and the mass of the SMBH, but the results have been inconclusive and sometimes controversial," Burke said.

The team compiled a large data set of actively feeding SMBHs to study the variability pattern of flickering. They identified a characteristic timescale, over which the pattern changes, that tightly correlates with the mass of the SMBH. The researchers then compared the results with accreting white dwarfs, the remnants of stars like our sun, and found that the same timescale-mass relation holds, even though [white dwarfs](#) are

millions to billions times less massive than SMBHs.



Explainer diagram - When Black Holes Line Up. Credit: Lucy Reading-Ikkanda / Simons Foundation

The light flickers are random fluctuations in a black hole's feeding process, the researchers said. Astronomers can quantify this flickering pattern by measuring the power of the variability as a function of timescales. For accreting SMBHs, the variability pattern changes from short timescales to long timescales. This transition of variability pattern happens at a characteristic timescale that is longer for more [massive black holes](#).

The team compared black hole feeding to our eating or drinking activity by equating this transition to a human belch. Babies frequently burp while drinking milk, while adults can hold in the burp for a more extended amount of time. Black holes kind of do the same thing while feeding, they said.

"These results suggest that the processes driving the flickering during accretion are universal, whether the central object is a supermassive black hole or a much more lightweight white dwarf," Shen said.

"The firm establishment of a connection between the observed light flicker and fundamental properties of the accretor will certainly help us better understand accretion processes," said Yan-Fei Jiang, a researcher at the Flatiron Institute and study co-author.

Astrophysical black holes come in a broad spectrum of mass and size. In between the population of stellar-mass black holes, which weigh less than several tens of times the mass of the sun, and SMBHs, there is a population of black holes called intermediate-mass black holes that weigh between about 100 and 100,000 times the mass of the sun.

IMBHs are expected to form in large numbers through the history of the universe, and they may provide the seeds necessary to grow into SMBHs later. However, observationally this population of IMBHs is surprisingly elusive. There is only one indisputably confirmed IMBH that weighs

about 150 times the mass of the sun. But that IMBH was serendipitously discovered by the gravitational wave radiation from the coalescence of two less-massive black holes.

"Now that there is a correlation between the flickering pattern and the [mass](#) of the central accreting object, we can use it to predict what the flickering signal from an IMBH might look like," Burke said.

Astronomers worldwide are waiting for the official kickoff of an era of massive surveys that monitor the dynamic and variable sky. The Vera C. Rubin Observatory in Chile's Legacy Survey of Space and Time will survey the sky over a decade and collect light flickering data for billions of objects, starting in late 2023.

"Mining the LSST data set to search for flickering patterns that are consistent with accreting IMBHs has the potential to discover and fully understand this long-sought mysterious population of black holes," said co-author Xin Liu, an astronomy professor at the U. of I.

This study is a collaboration with astronomy and physics professor Charles Gammie and astronomy postdoctoral researcher Qian Yang, the Illinois Center for Advanced Study of the Universe, and researchers at the University of California, Santa Barbara; the University of St. Andrews, U.K.; the Flatiron Institute; the University of Southampton, U.K.; the United States Naval Academy; and the University of Durham, U.K.

Burke, Shen and Liu also are affiliated with the Center for Astrophysical Surveys at the National Center for Supercomputing Applications at Illinois.

**More information:** A characteristic optical variability timescale in astrophysical accretion disks, *Science* (2021).



[science.sciencemag.org/lookup/.../1126/science.abg9933](https://science.sciencemag.org/lookup/.../1126/science.abg9933)

Provided by University of Illinois at Urbana-Champaign

Citation: Black hole size revealed by its eating pattern (2021, August 12) retrieved 9 April 2024 from <https://phys.org/news/2021-08-black-hole-size-revealed-pattern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.