

Black and white anglerfish hybridise producing viable descendants

August 2 2021



Anglerfish (*Lophius piscatorius*) is caught by researchers during monitoring cruise in the North Sea onboard German research vessel Walther Herwig III. Credit: [Randel Kreitsberg](#)/CC 4.0

Black and white anglerfish have always been considered to be two

separate species. Morphologically, they are mainly distinguished on the basis of the color of the peritoneum, the epithelium lining the intestinal cavity (black for black anglerfish and white for white anglerfish).

However, a new analysis led by the AZTI technology center, member of the BRTA, questions the effectiveness of this identification method and has also discovered the existence of thus far unknown hybrids, resulting from the mating between white and black anglerfish.

"We have shown for the first time that black and white anglerfish hybridize, and that these hybrids can reproduce and have viable offspring; furthermore, the percentage of hybrids in some areas is very high, up to 20 percent," explains marine genomics expert Naiara Rodríguez-Ezpeleta.

To carry out the study, developed within the framework of the GECKA project and published in the scientific journal *Evolutionary Applications*, AZTI researchers analyzed hundreds of anglerfish samples that were assigned to white or black anglerfish based on the color of the peritoneum. The samples were collected across the [species](#) distribution area in the Atlantic by collaborating institutes.

"When analyzing in the laboratory samples identified as white anglerfish, we realized that some of the specimens were genetically black anglerfish, so we came to the conclusion that the color of the peritoneum is not a reliable characteristic to assign species," explains the researcher. The study has also shown that there is hybridization between black and white anglerfish.

"We also saw that there are hybrids that arise from the mating between black and white anglerfish, but also between hybrids and white or black anglerfish, so we can ensure that hybrid anglerfish are viable and can reproduce, which could have consequences on the conservation of the species in the medium and long term, as it could lead to the

disappearance of the black and white anglerfish if the hybrids outperform them," adds Rodríguez-Ezpeleta.

Thus, it is crucial "to find out if hybridization has occurred recently and, although research has not yet been able to determine the reasons for it, it is possible that climate change has increased the coexistence of both species in the same area and, therefore, induced the existence of hybrids."

Impact on fisheries management

White anglerfish inhabit the Mediterranean Sea and the Northeast Atlantic, where they are managed by the International Council for the Exploration of the Seas (ICES) as three management units: North Shelf, North and South. Previous studies carried out to assess the population of this species in the Atlantic found no genetic differences between the management units, but this had to be confirmed through the analysis of a large number of genetic markers, such as in the study now developed by AZTI.

"By discarding hybrids and misassigned individuals, we focused on the white anglerfish and analyzed whether or not there were [genetic differences](#) between the management units. We found that the white anglerfish constitutes a single genetic population throughout the Atlantic, a crucial knowledge to inform [fisheries management](#) of this species and calculating Total Allowable Catch (TAC) recommendations," says the AZTI expert.

Therefore, the results obtained by AZTI are also relevant for the assessment of the biomass of white anglerfish and the management of the species, since the discovery opens up the possibility of managing the white anglerfish in the Atlantic as a single unit from now on, prioritizing biological characteristics over political and administrative considerations

for management unit definition, and thus providing greater accuracy when establishing the natural population of the species.

In addition, the study adds important information for the economy of the fishing sector and consumers, and it may have consequences on the price of black and white anglerfish, which is now different for each species.

"According to the data recorded by our fisheries assessment team, looking at the last two years, the price of white anglerfish in the first sale is usually about €4 or €5 per kilo, and that of black anglerfish, about €6 or €7. If you cannot visually distinguish both species, these prices shouldn't be different, either," adds the AZTI researcher.

The next objectives of the research, this time with the financial support of the General Secretariat of Fisheries of Spain, will be to study the impact of [hybrid anglerfish](#) on the evaluation and management of the species, including the economic impact of the potential decrease of the spawning biomass due to hybrids, and the species mixing.

More information: Imanol Aguirre-Sarabia et al, Evidence of stock connectivity, hybridization and misidentification in white anglerfish support the need of a genetics-informed fisheries management framework, *Evolutionary Applications* (2021). [DOI: 10.1111/eva.13278](https://doi.org/10.1111/eva.13278)

Provided by AZTI

Citation: Black and white anglerfish hybridise producing viable descendants (2021, August 2) retrieved 21 May 2024 from <https://phys.org/news/2021-08-black-white-anglerfish-hybridise-viable.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.