

Deforestation leads to flight loss in New Zealand insects

August 12 2021



A University of Otago study has revealed new evidence of rapid evolution in New Zealand's native insect species in response to increased exposure to wind from the loss of shelter due to deforestation.

The study, carried out by researchers in the Department of Zoology,

Stoneflies. Credit: Brodie Foster



found strong links between recent deforestation and loss of wings in <u>insect populations</u> in several southern New Zealand locations.

The research, led by Ph.D. student Brodie Foster and published in *Biology Letters*, used comparisons of insect populations around the alpine "treeline" in both forested and cleared sites, and found that recent loss of forest has increased the distribution of flightless insects which were formerly limited to alpine grassland ecosystems.

"Over a short period of just a few hundred years the study has found that many previously winged populations have evolved to become flightless," Mr Foster says.

The Otago team focussed on the endemic Zelandoperla stonefly species that frequently have flightless populations in mountainous regions, but typically have flighted populations in lower forested regions.

The researchers also found wingless populations occurring at much lower elevations in recently deforested areas. Genetic analysis revealed little difference between winged and flightless populations in these deforested areas, confirming that loss of flight has evolved very rapidly.

Study co-author Dr. Graham McCulloch, who led the <u>genetic analysis</u>, says due to the loss of forest cover, native insect species are increasingly vulnerable to exposure and have evolved flight-loss as a strategy to cope with these windy conditions.

Human expansion has drastically shifted the balance of many ecosystems across the globe, and how wild populations will continue to adapt and evolve to sudden changes such as deforestation remains largely unknown.

Traditionally, <u>evolutionary biologists</u> think about change occurring over



many millennia, whereas this study has found significant change taking place over the timescale of centuries.

However, Dr. McCulloch points out that while this study has shown the ability of these populations to adapt rapidly to <u>environmental changes</u>, it has also highlighted the increased vulnerability of these insects to local extinction as they become more isolated.

"While most of our flying insect species seem relatively secure, isolated flightless populations are potentially more vulnerable, as is also the case for our native birds."

The recent discovery of another flightless insect <u>population</u>, the Maungatua stonefly, in a very restricted region near Dunedin is an example of a vulnerable wild population.

Dr. McCulloch says it is essential that these unique, range-limited insects are conserved.

More information: Brodie J. Foster et al, Anthropogenic evolution in an insect wing polymorphism following widespread deforestation, *Biology Letters* (2021). DOI: 10.1098/rsbl.2021.0069

Provided by University of Otago

Citation: Deforestation leads to flight loss in New Zealand insects (2021, August 12) retrieved 12 July 2024 from <u>https://phys.org/news/2021-08-deforestation-flight-loss-zealand-insects.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.