

Fewer El Niño and La Niña events in a warmer world

August 26 2021



Surface ocean temperatures simulated at unprecedented resolution using a coupled atmosphere-ocean model. The extensive wavy cold structure in the equatorial Pacific corresponds to a tropical instability wave. Simulations were conducted on the IBS/ICCP supercomputer Aleph. Credit: Institute for Basic Science



The cycling between warm El Niño and cold La Niña conditions in the eastern Pacific (commonly referred to as the El Niño-Southern Oscillation, ENSO) has persisted without major interruptions for at least the last 11,000 years. This may change in the future according to a new study published in the journal *Nature Climate Change* by a team of scientists from the IBS Center for Climate Physics (ICCP) at Pusan National University in South Korea, the Max Planck Institute of Meteorology, Hamburg, Germany, and the University of Hawai'i at Mānoa, U.S.

The team conducted a series of global climate model simulations with an unprecedented spatial resolution of 10 km in the ocean and 25 km in the atmosphere. Boosted by the power of one of South Korea's fastest supercomputers (Aleph), the new ultra-high-resolution climate model simulations can now realistically simulate tropical cyclones in the atmosphere and tropical instability waves in the equatorial Pacific Ocean, which both play fundamental roles in the generation and termination of El Niño and La Niña events. "Our supercomputer ran non-stop for over one year to complete a series of century-long simulations covering present-day climate and two different global warming levels. The model generated 2 quadrillion bytes of data; enough to fill up about 2,000 hard disks," says Dr. Sun-Seon Lee who conducted the experiments.

Analyzing this enormous dataset, the team focused on a long-standing problem: how will ENSO change in response to increasing greenhouse gas concentrations. "Two generations of climate scientists have looked at this issue using climate models of varying complexity. Some models simulated weaker; others predicted larger eastern Pacific temperature swings in a future warmer climate. The jury was still out," says Prof. Axel Timmermann, co-corresponding author and Director of the ICCP.



He adds "What is common to these models is that their simulated temperatures in the equatorial Pacific, west of Galapagos, were always too cold compared to the observations. This prevented them from properly representing the delicate balance between positive and negative feedback processes that are important in the ENSO cycle."

By capturing small-scale climatic processes at the highest computationally possible resolution, the ICCP team was able to alleviate these ocean temperature biases, leading to substantial improvements in the representations of ENSO and its response to Global Warming. "The result from our computer simulations is clear: Increasing CO_2 concentrations will weaken the intensity of the ENSO temperature cycle," says Dr. Christian Wengel, first author of the study and former postdoctoral researcher at the ICCP, now at the Max Planck Institute of Meteorology in Hamburg in Germany.

By tracing the movement of heat in the coupled atmosphere/ocean system the scientists identified the main culprit of the collapse of the ENSO system: Future El Niño events will lose heat to the atmosphere more quickly due to the evaporation of water vapor, which has the tendency to cool the ocean. In addition, the reduced future temperature difference between the eastern and western tropical Pacific will also inhibit the development of temperature extremes during the ENSO cycle. However, these two factors are partly offset by a projected future weakening of tropical instability waves. Normally these oceanic waves, which can encompass up to 30% of the earth's entire circumference, develop during La Niña conditions. They replace colder equatorial waters with warmer off-equatorial water, thereby accelerating the demise of a La Niña event. The new <u>computer simulations</u>, which resolve the detailed structure of these waves, demonstrate that the associated negative feedback for ENSO will weaken in the future.

"There is a tug-of-war between positive and negative feedbacks in the



ENSO system, which tips over to the negative side in a warmer climate. This means future El Niño and La Niña events cannot develop their full amplitude anymore," says Prof. Malte Stuecker, co-author of the study and now assistant professor at the Department of Oceanography and the International Pacific Research Center at the University of Hawai'i at Mānoa.

Even though the year-to-year fluctuations in eastern equatorial Pacific temperatures are likely to weaken with human-induced warming according to this new study, the corresponding changes in El Niño and La Niña-related rainfall extremes will continue to increase due to an intensified hydrological cycle in a warmer climate, as shown in recent studies by scientists from the ICCP and their international collaborators.

"Our research documents that unabated warming is likely to silence the world's most powerful natural <u>climate</u> swing which has been operating for thousands of years. We don't yet know the ecological consequences of this potential no-analog situation," says Axel Timmermann. "But we are eager to find out."

More information: Future high-resolution El Niño/Southern Oscillation dynamics, *Nature Climate Change* (2021). DOI: <u>10.1038/s41558-021-01132-4</u>, <u>www.nature.com/articles/s41558-021-01132-4</u>

Provided by Institute for Basic Science

Citation: Fewer El Niño and La Niña events in a warmer world (2021, August 26) retrieved 3 May 2024 from <u>https://phys.org/news/2021-08-el-nio-la-nia-events.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.