

Small satellites have a role in assessing atmospheric particulates, research shows

August 18 2021



Credit: Pixabay/CC0 Public Domain

Small commercial satellites have a role in atmospheric particulates investigations, according to recently published thesis research by a research associate with the Earth System Science Center at The

University of Alabama in Huntsville (UAH), a part of the University of Alabama System.

The research by Jeanné le Roux (M.S., Earth System Science, 2020) is among the first to use commercial high resolution small [satellite](#) (smallsat) data for air quality investigations.

The peer-reviewed paper appeared in *Remote Sensing* and was co-authored by le Roux, Dr. Sundar Christopher, UAH professor of atmospheric and Earth science, and Manil Maskey, a researcher at NASA's Marshall Space Flight Center. The research was supported by a NASA grant for its Commercial Smallsat Data Acquisition program.

Le Roux looked at applying the data from PlanetScope sensors aboard 100-plus Dove smallsats deployed by the company Planet for use in fine particulate matter air quality assessments. Dove satellites are smallsats in 3U configurations, which basically means each is the size of three cube satellites (CubeSats) joined together, measuring 10 x 10 x 30 centimeters.

"This was an initial investigation into the feasibility of using commercial high resolution CubeSat data," says le Roux, who is from Riverside, Calif., and also works as a research associate with NASA's Interagency Implementation and Advanced Concepts Team (IMPACT).

"While this data shows a lot of promise for air quality applications, there are certain considerations that need to be taken into account, primarily related to inconsistencies that can arise when data from 130-plus individual sensors are combined."

Planet was the first in the industry to launch such a large number of smallsats in the same network for the specific purpose of Earth observation, but there are many other companies that have or are

planning smallsat networks for various purposes.

Atmospheric scientists use various satellite data to augment a large network of ground-based air quality monitoring stations located in the U.S. and around the globe, such as the Environmental Protection Agency's Air Quality System.

"The issue with these ground networks is that while they take frequent measurements, they only provide air quality information at the specific point location of the ground monitoring site," le Roux says. "Satellite data can be used to help fill in the gaps and estimate air quality parameters over vast and continuous spatial scales."

Commercial enterprises have been at the forefront of providing very high-resolution satellite imagery available for purchase, le Roux says.

"Until government-funded satellites are designed to compete, this commercial data will be highly sought after for a very wide variety of scientific applications," she says.

"Data from the company Planet specifically is unique, in that it provides high resolution coverage of most of the Earth's landmass on a daily basis. Most other commercial high-resolution satellites do not offer global coverage at this frequency, and are rather tasked to focus on specific areas of the Earth based on customer requests."

While there are government satellites that provide air quality information, none of them have been developed to do so at a spatial resolution that is as detailed as that currently offered by commercial satellites, le Roux says.

"For example, NASA's Moderate Resolution Imaging Spectroradiometer (MODIS) can be used to derive various air quality parameters with a

pixel size as low as 250 meters, but PlanetScope's pixel size is still considerably smaller at only 3 meters," le Roux says.

While it is possible that the government will invest in technology similar to Planet's commercial network in the future, to date commercial enterprises have been the pathfinders for this approach to collecting remotely sensed imagery, she says.

Le Roux did find some drawbacks to using smallsats for atmospheric particulate research.

"The main drawback to this data is potential consistency issues that arise, since data are combined from over 100 individual sensors," she says.

"While great care is taken to make imagery from the various sensors as consistent as possible, geometric and radiometric calibration quality concerns have been brought up in past studies."

Also, CubeSats have a limited number of bands to view the Earth, she says. While the MODIS has 36 channels, commercial [sensors](#) have three to four bands, typically in the visible to the near-infrared part of the spectrum. That's a drawback for studies that need more coverage in the electromagnetic spectrum.

"One of the first things that we need to understand is the quality of the measurements using calibration concepts. Then we need to assess the geolocation accuracies," le Roux says. "Our work provides the framework and methods for assessing these accuracies."

The technology employed in Planet's constellation of smallsats continues to improve so the quality of data should continue to be evaluated in future studies, le Roux says.

"The benefits of having this frequent high-resolution imagery often

outweigh some of the data quality concerns."

More information: Jeanné le Roux et al, Exploring the Use of PlanetScope Data for Particulate Matter Air Quality Research, *Remote Sensing* (2021). [DOI: 10.3390/rs13152981](https://doi.org/10.3390/rs13152981)

Provided by University of Alabama in Huntsville

Citation: Small satellites have a role in assessing atmospheric particulates, research shows (2021, August 18) retrieved 2 February 2023 from <https://phys.org/news/2021-08-small-satellites-role-atmospheric-particulates.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.