

## New solvents to break down plant cellulose for bioethanol

August 10 2021



Suitable solvents are needed to produce ethanol from cellulosic plant biomass. We have found that liquid zwitterions are suitable for it, but it has been difficult to liquefy zwitterions and the species have been strictly limited. Therefore, the solid zwitterions were liquefied through making deep eutectic solvents, which are liquids from solid-solid mixtures. The liquefied zwitterionic solvent was able to dissolve the cellulose. Credit: Kanazawa University

Scientists from the Institute of Science and Engineering at Kanazawa University have developed new solvent mixtures to break down the tough structure of plant cellulose for the production of bioethanol. These new solvents work under mild conditions, have reduced toxicity and are more environmentally friendly compared with currently available



solvents. This work may lead to improved technologies for the conversion of currently unused biomass to fuel.

Biofuels, such as ethanol produced from switchgrass or sugarcane, may allow us to reduce our reliance on nonrenewable fossil fuels. However, the process of making biofuels usually requires breaking down the <u>cellulose</u> from plants, which consists of long polymer chains, into smaller sugar <u>molecules</u>. This is not a simple task, as illustrated by how unappetizing tree bark or grass stalks look to us humans. Cellulose has a complex hydrogen bonding network, which makes it extremely chemical resistant. Current methods for processing cellulose rely on harsh reaction conditions and <u>toxic chemicals</u>.

Now, scientists from Kanazawa University have used a special class of molecules called "zwitterions" to create novel solvents with the ability to dissolve cellulose. Zwitterions are special in that they have both a positive and <u>negative charge</u>, but at different locations on the molecule so they cannot neutralize each other. These charges are highly effective at disrupting the <u>hydrogen bonds</u> keeping the cellulose from being broken down. "Because almost all zwitterions are solid under normal reaction conditions, our experiments used eutectic mixtures," first author Gyanendra Sharma explains. A eutectic system is a mixture of substances with a melting point lower than that of its constituent parts. This is accomplished by using molecules with different structures, so that regular crystals are harder to form.

In these experiments, the team mixed four different zwitterions at various ratios. They found 22 combinations that were liquid below 100°C. Of these, two mixtures were also highly effective at dissolving cellulose. "Our work shows that it is possible to replace many of the toxic chemicals used today with more environmentally friendly alternatives as we move towards a more renewable energy ecosystem," senior author Kosuke Kuroda says. This research demonstrates the



potential of using combinations of zwitterions to create mixtures with properties not possessed by any molecule individually.

The work was published in Carbohydrate Polymers.

**More information:** Gyanendra Sharma et al, Polar zwitterion/saccharide-based deep eutectic solvents for cellulose processing, *Carbohydrate Polymers* (2021). DOI: 10.1016/j.carbpol.2021.118171

Provided by Kanazawa University

Citation: New solvents to break down plant cellulose for bioethanol (2021, August 10) retrieved 27 June 2024 from <u>https://phys.org/news/2021-08-solvents-cellulose-bioethanol.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.