

Unveiling a century-old mystery: Where the Milky Way's cosmic rays come from

August 23 2021




Figure 1. Schematic images of gamma-ray production from cosmic-ray protons and electrons. Cosmic-ray protons interact with interstellar protons such as molecular and atomic hydrogen gas. The interaction creates neutral pion that quickly decays into two gamma-ray photons (hadronic process). Cosmic-ray electrons energize interstellar photons (mainly Cosmic Microwave Background; CMB) into gamma-ray energy via inverse Compton scattering (leptonic process). Credit: Astrophysics Laboratory, Nagoya University

Astronomers have succeeded for the first time in quantifying the proton and electron components of cosmic rays in a supernova remnant. At least 70% of the very-high-energy gamma rays emitted from cosmic rays are due to relativistic protons, according to the novel imaging analysis of radio, X-ray, and gamma-ray radiation. The acceleration site of protons, the main components of cosmic rays, has been a 100-year mystery in modern astrophysics, this is the first time that the amount of cosmic rays being produced in a supernova remnant has been quantitatively shown and is an epoch-making step in the elucidation of the origin of cosmic

rays.

The origin of cosmic rays, the particles with the highest energy in the universe, has been a great mystery since their discovery in 1912. Because cosmic rays promote the chemical evolution of interstellar matter, understanding their origin is critical in understanding the evolution of our Galaxy. The cosmic rays are thought to be accelerated by [supernova remnants](#) (the after-effects of supernova explosions) in our Galaxy and traveled to the Earth at almost the speed of light. Recent progress in gamma-ray observations has revealed that many supernova remnants emit [gamma-rays](#) at teraelectronvolts (TeV) energies. If gamma rays are produced by protons, which are the main component of cosmic rays, then the supernova remnant origin of cosmic rays can be verified. However, gamma rays are also produced by electrons, it is necessary to determine whether the [proton](#) or electron origin is dominant, and to measure the ratio of the two contributions (see also Figure 1). The results of this study provide compelling evidence of gamma rays originating from the proton component, which is the main component of cosmic rays, and clarify that Galactic cosmic rays are produced by supernova remnants.

The originality of this research is that [gamma-ray radiation](#) is represented by a linear combination of proton and electron components. Astronomers knew a relation that the intensity of gamma-ray from protons is proportional to the interstellar gas density obtained by radio-line imaging observations. On the other hand, gamma-rays from electrons are also expected to be proportional to X-ray intensity from electrons. Therefore, they expressed the total gamma-ray intensity as the sum of two gamma-ray components, one from the proton origin and the other from the electron origin. This led to a unified understanding of three independent observables (Figure 2). This method was first proposed in this study. As a result, it was shown that gamma rays from protons and electrons account for 70% and 30% of the total gamma-rays,

respectively. This is the first time that the two origins have been quantified. The results also demonstrate that gamma rays from protons are dominated in interstellar gas-rich regions, whereas gamma rays from electrons are enhanced in the gas-poor region. This confirms that the two mechanisms work together and supporting the predictions of previous theoretical studies.




Figure 2. Maps of gamma-ray intensity N_g , interstellar gas density N_p , and X-ray intensity N_x . Credit: Astrophysics Laboratory, Nagoya University

"This novel method could not have been accomplished without international collaborations," says Emeritus Professor Yasuo Fukui at Nagoya University. He led this project and has accurately quantified interstellar gas density distribution using the NANTEN radio telescope and Australia Telescope Compact Array since 2003. Although the gamma ray resolution was insufficient to perform a full analysis at that time, Professor Gavin Rowell and Dr. Sabrina Einecke of the University of Adelaide and the H.E.S.S. team dramatically improved the spatial resolution and sensitivity of gamma rays over the years, making it possible to compare them precisely with interstellar gas. Dr. Hidetoshi

Sano of the National Astronomical Observatory of Japan led the X-ray imaging analysis of archival datasets from the European X-ray satellite XMM-Newton. Dr. Einecke and Prof. Rowell worked closely with Prof. Fukui and Dr. Sano on making the detailed studies that examined the correlations across the gamma-ray, X-ray and radio emission. "This novel method will be applied to more [supernova remnants](#) using the next-generation [gamma-ray](#) telescope CTA (Cherenkov Telescope Array) in addition to the existing observatories, which will greatly advance the study of the [origin](#) of [cosmic rays](#)."




Figure 3. Three-dimensional fitting of a flat plane expressed by an equation of $N_g = a N_p + b N_x$, where a and b are constants. The data points are colored by the code in the figure according to N_g and are shown by filled and open symbols for those above and below the plane. The blue, green, yellow, and red represent N_g is less than 1.2 counts arcmin⁻², 1.2–1.7 counts arcmin⁻², 1.7–2.2 counts arcmin⁻², and greater than 2.2 counts arcmin⁻², respectively. The blue, green, orange, red, and purple dashed lines on the best-fit plane indicate 1.0, 1.5, 2.0, 2.5, and 3.0 counts arcmin⁻², respectively. Credit: Astrophysics Laboratory, Nagoya University

More information: Yasuo Fukui et al, Pursuing the Origin of the Gamma Rays in RX J1713.7-3946 Quantifying the Hadronic and Leptonic Components, *The Astrophysical Journal* (2021). [DOI: 10.3847/1538-4357/abff4a](https://doi.org/10.3847/1538-4357/abff4a)

Provided by Nagoya University

Citation: Unveiling a century-old mystery: Where the Milky Way's cosmic rays come from (2021, August 23) retrieved 29 January 2023 from <https://phys.org/news/2021-08-unveiling-century-old-mystery-milky-cosmic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.