

Ways to improve magpie goose management on mango farms

August 11 2021



CDU PhD student Amélie Corriveau is researching ways to help reduce the damage made by magpie geese on Northern Territory farms. Credit: Charles Darwin University

Research from Charles Darwin University (CDU) has identified ways to improve management of magpie geese to better assist mango growers as the Northern Territory goes into mango season.

Ph.D. student and environmental science researcher Amélie Corriveau has recently completed her research in studying the movements and diet



of the magpie goose, a large and iconic bird most abundant in the Top End of the Northern Territory.

Although protected wildlife and a totemic animal for Indigenous people, magpie <u>geese</u> can be a headache to local farmers as they feed on crops and cause financial losses.

Ms Corriveau investigated the seasonal movements and food resources of magpie geese in the Northern Territory to help farmers identify appropriate approaches to deal with these birds.

"My research aimed to understand where geese coming on farms were from, whether the same birds came back to the region every year, and how they used the resources available in the agricultural landscape," Ms Corriveau said.

"The findings showed that the birds were opportunistic and adaptable, but also that they used <u>prior knowledge</u>—the same geese did tend to come back every year. That means that geese are going to be an ongoing challenge from growers, as they can easily locate attractive resources and go back to them."

Magpie geese were also found to be quite flexible in the areas they use and could go from one local area to another on a whim.

"They could relocate easily and find resources within short distances. They fed on both agricultural and natural occurring plants, but their diet varied among individuals, sites and weeks of the season," Ms Corriveau said.

These findings provide new insights into magpie goose behavior and habitat use, which can help farmers manage them.



"They seem to establish within areas where they can have access to shade, water and freshly irrigated grass, which are very attractive in the late dry season."

This means farmers could reduce the attractiveness of their farms for geese by using underground irrigation systems or night-time irrigation schedules, minimizing available water on farms or removing fruit falls.

Setting up alternative areas to divert the birds away from farms could also help reduce damage by keeping <u>birds</u> chased from farms away for longer.

"This is a very complex issue and there is no magical solution. Systemic damage assessments are needed to quantify the actual damage made by geese" Ms Corriveau said.

"Management needs to be adaptable to how the species behave from one season to another. Farmers could also pool resources and coordinate among neighbors to maximize the effectiveness of management actions on farms."

Ms Corriveau also said the magpie goose is a protected species and we needed to ensure its long-term conservation.

"The population of <u>magpie</u> geese can fluctuate widely, and the geese have a very high socio-economic value in the Northern Territory," she said.

"They bring in tourism for hunting and bird watching, so it is important to find approaches that enable both the conservation of the species and development of agriculture in northern Australia."



Provided by Charles Darwin University

Citation: Ways to improve magpie goose management on mango farms (2021, August 11) retrieved 27 June 2024 from <u>https://phys.org/news/2021-08-ways-magpie-goose-mango-farms.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.