

## Risk of airborne transmission of avian influenza from wild waterfowl to poultry negligible

September 28 2021



Credit: Pixabay/CC0 Public Domain

Research by Wageningen Bioveterinary Research (WBVR) has shown that the risk of airborne transmission of high pathogenic avian influenza



virus from infected wild birds is negligible. The research looked specifically at the airborne movement of particles from wild waterfowl droppings in the vicinity of poultry farms during the risk season for avian influenza (October to March). It also considered transmission via aerosolization, with the exhalations or coughs of wild waterfowl infected with avian influenza virus finding their way into the ventilation systems of poultry farms. As a precaution, it's important that the carcasses of wild waterfowl or other wild birds that have died of high pathogenic avian influenza are removed from their habitat as soon as possible. If not, scavengers eating the carcasses could cause feathers to become distributed. Feathers of wild birds that died of, and if the wild bird died of high pathogenic avian influenza contain the virus, which can then the virus can survive for a long time in those feathers.

## Introduction of the virus to poultry

"Wild waterfowl such as ducks, geese, swans and gulls are natural reservoirs of avian influenza viruses. These wild waterfowl can play an important role in introducing the virus to poultry, because they shed avian influenza virus when they're infected. Traces of the droppings of wild <u>birds</u> infected with avian influenza, when deposited in the vicinity of poultry farms, can find their way into those farms. For example, they can become attached to boots, clothing or tools, or they can be carried in with bedding material stored outside the poultry houses, or they can be brought in through rats and mice carrying the droppings on their fur or legs.

Furthermore, previous experiences have suggested that the avian influenza virus can be transported from one farm to another if it is produced in large quantities by large numbers of infected chickens in a poultry house. This project investigated whether there is a risk of the avian influenza virus being transmitted into poultry houses from infected wild birds in the vicinity of those poultry houses," says Armin Elbers,



project leader for the research and a senior epidemiologist at WBVR.

## **Risk analysis**

The focus of this risk analysis of airborne transmission of high pathogenic avian influenza virus derived from the contaminated droppings of wild waterfowl or from an aerosol produced by the exhalations or coughs of wild waterfowl infected with avian influenza virus was selected because the risk of such transmission routes might possibly be mitigated through the use of windbreak mesh.

The research identified all the relevant building blocks for carrying out a semi-quantitative risk analysis by means of an extensive literature review, and executing a field experiment to investigate the potential of airborne movement of bird dropping particles during the high-risk avian influenza season.

Provided by Wageningen University

Citation: Risk of airborne transmission of avian influenza from wild waterfowl to poultry negligible (2021, September 28) retrieved 30 April 2024 from <u>https://phys.org/news/2021-09-airborne-transmission-avian-influenza-wild.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.