

# Decaying forest wood releases 10.9 billion tons of carbon yearly, which will increase with climate change

September 2 2021, by Marisa Stone



Credit: Pixabay/CC0 Public Domain

If you've wandered through a forest, you've probably dodged dead, rotting branches or stumps scattered on the ground. This is "deadwood,"



and it plays several vital roles in forest ecosystems.

It provides habitat for small mammals, birds, amphibians and insects. And as deadwood decomposes it <u>contributes to the ecosystem's cycle of</u> <u>nutrients</u>, which is important for plant growth.

But there's another important role we have little understanding of on a global scale: the <u>carbon</u> deadwood releases as it decomposes, with part of it going into the soil and part into the atmosphere. Insects, such as termites and wood borers, can accelerate this process.

The world's deadwood currently stores <u>73 billion tons</u> of carbon. <u>Our</u> <u>new research in Nature</u> has, for the first time, calculated that 10.9 billion tons of this (around 15%) is released into the atmosphere and soil each year—<u>a little more than</u> the world's emissions from burning fossil fuels.

But this amount can change depending on insect activity, and will likely increase under <u>climate</u> change. It's vital deadwood is considered explicitly in all future climate change projections.

## An extraordinary, global effort

Forests are crucial carbon sinks, where living trees capture and store carbon dioxide from the atmosphere, helping to regulate climate. Deadwood—including fallen or still-standing trees, branches and stumps—makes up 8% of this carbon stock in the world's forests.

Our aim was to measure the influence of climate and insects on the rate of decomposition—but it wasn't easy. Our <u>research paper</u> is the result of an extraordinary effort to co-ordinate a large-scale cross-continent field experiment. More than 30 research groups worldwide took part.

Wood from more than 140 tree species was laid out for up to three years



at 55 forest sites on six continents, from the Amazon rainforest to Brisbane, Australia.Half of these wood samples were in closed mesh cages to exclude insects from the decomposition process to test their effect, too.

Some sites had to be protected from elephants, another was lost to fire and another had to be rebuilt after a flood.

#### What we found

Our research showed the rate of deadwood decay and how insects contribute to it depend very strongly on climate.

We found the rate increased primarily with rising temperature, and was disproportionately greater in the tropics compared to all other cooler climatic regions.

In fact, deadwood in <u>tropical regions</u> lost a median mass of 28.2% every year. In cooler, temperate regions, the median mass lost was just 6.3%.

More deadwood decay occurs in the tropics because the region has greater biodiversity (more insects and fungi) to facilitate decomposition. As insects consume the wood, they render it to small particles, which speed up decay. The insects also <u>introduce fungal species</u>, which then finish the job.

Of the 10.9 billion tons of carbon dioxide released by deadwood each year, we estimate insect activity is responsible for 3.2 billion tons, or 29%.

Let's break this down by region. In the tropics, insects were responsible for almost one-third of the carbon released from deadwood. In regions with low temperatures in forests of northern and temperate



latitudes—such as in Canada and Finland—insects had little effect.

### What does this mean in a changing climate?

<u>Insects are sensitive to climate change</u> and, with <u>recent declines in insect</u> <u>biodiversity</u>, the current and future roles of insects in deadwood are uncertain.

But given the vast majority of deadwood decay occurs in the tropics (93%), and that this region in general is set to become even warmer and wetter under climate change, it's safe to say climate change will increase the amount of carbon deadwood releases each year.



Credit: AI-generated image (disclaimer)



It's also worth bearing in mind that the amount of carbon dioxide released is still only a fraction of the total annual global deadwood carbon stock. That is, 85% of the global deadwood carbon stock remains on forest floors and continues to store carbon each year.

We recommend deadwood is left in place—in the forest. Removing deadwood may not only be <u>destructive for biodiversity and the ability of forests to regenerate</u>, but it could actually substantially increase atmospheric carbon.

For example, if we used deadwood as a biofuel it could release the carbon that would otherwise have remained locked up each year. If the world's deadwood was removed and burned, it would be release eight times more carbon than what's currently emitted from burning fossil fuels.

This is particularly important in cooler climatic regions, where decomposition is slower and deadwood remains for several years as a vital carbon sink.

#### What next?

The complex interplay of interactions between insects and climate on deadwood carbon release makes future climate projections a bit tricky.

To improve climate change predictions, we need much more detailed research on how communities of decomposer <u>insects</u> (such as the numbers of individuals and species) influence deadwood decomposition, not to mention potential effects from insect diversity loss.

But insect diversity loss is also likely to vary regionally and would require long-term studies over decades to determine.



For now, climate scientists must take the enormous annual emissions from deadwood into account in their research, so humanity can have a better understanding of <u>climate change</u>'s cascading effects.

**More information:** Sebastian Seibold et al, The contribution of insects to global forest deadwood decomposition, *Nature* (2021). <u>DOI:</u> <u>10.1038/s41586-021-03740-8</u>

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Decaying forest wood releases 10.9 billion tons of carbon yearly, which will increase with climate change (2021, September 2) retrieved 21 May 2024 from https://phys.org/news/2021-09-forest-wood-billion-tons-carbon.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.