

Hot-band absorption of Indocyanine Green for advanced anti-Stokes fluorescence bioimaging

September 15 2021



a, The molecular structure of ICG. b, Schematic illustration of energy level transition in HBA ASF. The thermal activation (red arrow) acts on the molecule in the ground state with lower energy at first and raises the molecule to a higher vibrational level of the ground state, then the molecule absorbs a long-wavelength photon (pink arrow) to reach the excited state, and finally emits a shorter-wavelength photon (orange arrow). c, Temperature dependence of ICG's



ASF spectra (800 nm – 900 nm) excited by a 915 nm CW laser. d, Fluorescence images of a tumor mouse reflecting the enhancement of ASF intensity of ICG in the breast tumor as the temperature rises. Credit: by Jing Zhou, Xiaoxiao Fan, Di Wu, Jie Liu, Yuhuang Zhang, Zikang Ye, Dingwei Xue, Mubin He, Liang Zhu, Zhe Feng, Andrey N. Kuzmin, Wen Liu, Paras N. Prasad, Jun Qian

Hot-band absorption based anti-Stokes fluorescence (ASF) of Indocyanine Green (ICG), much brighter than luminescence from rareearth ion doped nanoparticles, was observed and studied by scientists in China and U.S. recently. They found that the ASF of ICG can be used for tomography and blood flow velocity measurement of cerebral vessels, reflecting the change of temperature, and achieving multi-organ simultaneous imaging, and has considerable application prospects in biological imaging, sensing and even clinical translation.

There are four common types of anti-Stokes fluorescence (ASF): (i) direct multiphoton absorption (MPA) process, (ii) upconversion (UC) process based on multistep absorption through intermediate energy levels, (iii) thermally activated delayed fluorescence (TADF) process, and (iv) hot-band absorption (HBA) process. The occurrence of MPA fluorescence generally requires extremely high excitation intensity and is usually achieved by using expensive femto- or pico-second pulsed lasers. UC processes in rare-earth ion doped nanoparticles (UCNPs), or triplettriplet annihilation (TTA) based UC, can be obtained by using inexpensive continuous wave (CW) diode lasers. However, the absorption cross section of UCNPs is relatively small, resulting in low UC efficiency. TTA based metal complexes/organic compounds systems have larger absorption and higher quantum efficiency to be more efficient upconverters than UCNPs. Unfortunately, photostability of TTA based upconverters is relatively low due to strong quenching processes caused by molecular oxygen. TADF and HBA process in



organic molecules excited by the CW laser are attractive anti-Stokes processes. What's more, the potential of them to provide information about temperature in excited volume, makes them more attractive for application in bioimaging.

In a new paper published in *Light Science & Application*, a team of scientists, led by Professor Jun Qian from State Key Laboratory of Modern Optical Instrumentations, College of Optical Science and Engineering, Zhejiang University, China, and Professor Paras N Prasad from Institute for Lasers, Photonics and Biophotonics, State University of New York at Buffalo, U.S., discovered and studied the HBA based ASF in Food and Drug Administration (FDA)-approved Indocyanine Green (ICG). Based on thermal sensitivity, they applied ICG's ASF for evaluating the thermal state of subcutaneous tumors of mice during photothermal treatment. Besides, ICG's ASF is much stronger than typical UC fluorescence in UCNPs excited at 980 nm, with negligible thermal damage to biological tissues. Deep volume tomography of cerebral blood vessels and measurement of blood flow velocity of mice were performed by using the ASF of ICG. Moreover, in combination with L1057 nanoparticles (NPs), which absorb the ASF of ICG and emit beyond 1100 nm, these two probes generate multi-mode images in two fluorescent channels under the excitation of a single 915 nm CW laser. One channel is used to monitor two overlapping organs, urinary system and blood vessel of the rat, while the other shows urinary system only.





a, Photothermal effects on rat's hind limb and liver irradiating with the 915 nm or 980 nm CW laser. b, In vivo ASF wide-field imaging of biliary tracts of two rats after receiving injection of NaYF4: Yb3+, Tm3+ and ICG respectively. c, In vivo ASF wide-field microscopic image of brain blood vessels (depth at 250 μ m)



of the mouse injected with ICG. d, Flow velocity measurements of three sampled brain blood vessels. Credit: by Jing Zhou, Xiaoxiao Fan, Di Wu, Jie Liu, Yuhuang Zhang, Zikang Ye, Dingwei Xue, Mubin He, Liang Zhu, Zhe Feng, Andrey N. Kuzmin, Wen Liu, Paras N. Prasad, Jun Qian

In the HBA process, electrons in ICG molecules absorb photons from the upper, thermally populated, vibrational levels of ground state. The excitation decays to the lower vibration levels of the ground state, thus emitting photons with higher energy compared to that of absorbed initially. These scientists summarize the main work of HBA based ASF of ICG:

"We discovered bright ASF in ICG under the excitation of a 915 nm CW laser, and compared it with ASF of UCNPs excited by CW laser. The result is that the ASF of ICG is much brighter than that of UCNPs, which strongly attracted our attention. To find out the generation mechanism of ICG's ASF, we carried out rigorous verification experiments and concluded its generation mechanism to be HBA. Then we explored its possibility in thermal sensing applications. Evaluation of the thermal state of subcutaneous tumors during photothermal treatment and high temperature indication were achieved by using ICG's ASF. Cerebral blood vessel tomography and blood stream velocimetry of mice were performed too. Considering the surgical operation, we proved a concept of real-time in vivo multi-mode imaging enabling high contrast and selective detection of adjacent organs (urinary system and blood vessels) by combining ICG with fluorescent organic polymer dots L1057 under a single 915 nm CW laser excitation. This new imaging technique can be helpful for intraoperative real-time monitoring and avoiding accidental surgery injury."





a, The optical principle of multi-mode imaging. 915 nm is the excitation wavelength for ICG to produce ASF, and is also the absorption peak of L1057 NPs, and the ASF of ICG in channel 1 is exactly located in the high absorption spectral region of L1057 NPs. Meanwhile Stokes fluorescence (SF) signals of L1057 NPs in channel 2 do not have crosstalk with that of ICG. b, Injection mode of multi-mode imaging of urinary system and blood vessels. ICG is injected into the blood vessels and L1057 NPs are injected into the urinary system. c, Multi-mode pseudo color imaging of ureters and blood vessels of the rat treated with ICG and L1057 NPs under the excitation of a 915 nm CW laser. In channel 1, the blood vessels present bright positive visualization (displayed in pseudo color green), and the ureters which absorb the background ASF of ICG present obvious dark negative visualization (displayed in pseudo color pink), achieving automatic two-organ simultaneous visualization. Meanwhile, in channel 2, only ureters have signals (displayed in pseudo color pink), achieving the single-target visualization. Credit: by Jing Zhou, Xiaoxiao Fan, Di Wu, Jie Liu, Yuhuang Zhang, Zikang Ye, Dingwei Xue, Mubin He, Liang Zhu, Zhe



Feng, Andrey N. Kuzmin, Wen Liu, Paras N. Prasad, Jun Qian

"Interestingly, we found that ICG could generate visible ASF (

"We expect ASF of ICG to be more deeply and widely used and other fluorophores with HBA induced ASF features can be synthesized for bioimaging, sensing, theranostics and infrared light perception in the future." the scientists forecast.

More information: Jing Zhou et al, Hot-band absorption of indocyanine green for advanced anti-stokes fluorescence bioimaging, *Light: Science & Applications* (2021). DOI: 10.1038/s41377-021-00627-1

Provided by Chinese Academy of Sciences

Citation: Hot-band absorption of Indocyanine Green for advanced anti-Stokes fluorescence bioimaging (2021, September 15) retrieved 28 April 2024 from <u>https://phys.org/news/2021-09-hot-band-absorption-indocyanine-green-advanced.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.