


Physicist observes the first unpaired Weyl magnetic monopole

September 14 2021



The 37 unpaired Weyl points in the 3D reciprocal space of GaPt, and the Berry curvature field (analogous to magnetic field) component in the $kz=0$ plane.
 Credit: Dr. Ma Junzhang

Similar to a magnet that always has both south and north poles, a kind of special quasiparticles in condensed matter called "Weyl Fermions" always appear in pairs with opposite chirality. There had been no

experimental report that unpaired Weyl points exist in condensed matter until recently, a City University of Hong Kong (CityU) physicist observed the first unpaired singular Weyl magnetic monopole in a specific kind of single crystalline solid.

The research work was co-led by Dr. Ma Junzhang, Assistant Professor in the Department of Physics at CityU and collaborating scientists from École Polytechnique Fédérale de Lausanne (EPFL) and Paul Scherrer Institute (PSI) in Switzerland. It was published in *Nature Communications*, titled "Observation of a singular Weyl point surrounded by charged nodal walls in PtGa."

"Our group is the first to see and record an unpaired Weyl [magnetic monopole](#) in the experiment. The findings open a new avenue to search for the bulk topological properties of unpaired Weyl magnetic monopoles in solids, which will promote the understanding of basic topological physics, and the application of Weyl semimetals into spintronics," Dr. Ma said.


Weyl points: similar properties with magnetic monopoles

In the real world, a magnet must have south and north poles simultaneously. Modern physics theories suggest the existence of a magnetic [monopole](#), i.e. a quantum elementary unit of magnet with just one pole. But so far, there has been no known experimental or observational evidence that magnetic monopoles exist. Searching for magnetic monopole has been a dream for the modern physics community.

Similarly, Weyl points in condensed matter (including semimetal crystal) have the similar properties as the magnetic monopoles. "Therefore, Weyl

points in condensed matter are also called Weyl magnetic monopoles," explained Dr. Ma.

But there is one difference for Weyl points, based on many physicists' perceptions. It is widely believed that the existence of an unpaired Weyl magnetic monopole in solids is impossible due to the well-known Nielsen-Ninomiya no-go theorem. As a result, it is thought that Weyl magnetic monopoles in semimetals should always appear in pairs with opposite charges in the 3D momentum space. And the projections of the monopoles on the surface of the single crystal should always be connected by one kind of conducting state called Fermi arcs, which acts like an electronic channel that electrons can transport through it.



Experimental observation of unpaired magnetic monopole in energy-momentum space of PtGa. Credit: Dr. Ma Junzhang

No successful experimental precedent

Considerable theoretical efforts have been dedicated to searching unpaired Weyl monopoles beyond this no-go theorem, but no successful experiment was reported in the past. Many scientists thought that such kind of "unpaired substance" was hard to exist in single crystalline solids.

But Dr. Ma didn't think so. He searched thousands of compounds in the database and finally discovered that a few dozens of them are the [potential candidates](#) hosting unpaired Weyl monopoles. His suggestions were supported by theoretical colleagues Dr. Wu Quansheng, and Professor Oleg Yazyev from EPFL. After that, a team led by Dr. Ma and Professor Shi Ming from PSI started a series of photoemission experiments at the Swiss Light Source at PSI and successfully proved the primary ideas.

Combining angle-resolved photoemission spectroscopy (ARPES) experiments with density functional theory calculations, the research team revealed for the first time that unpaired Weyl monopoles can emerge in various compounds. In studying the promising candidate, semimetal crystal platinum gallium (PtGa), the research team identified 37 Weyl monopoles in the momentum space, and found out that the system hosts odd number 37 Weyl points with either +1 charge or -1 charge that prevent them pair together.

They also demonstrated experimentally for the first time that unpaired magnetic monopole can exist in the momentum space of solid without surface Fermi arc. The geometry of the unpaired Weyl point is quite different from that of normal Weyl semimetals, so the properties are expected to change significantly in this new kind of materials, which defines a new study direction.

A new research direction

Dr. Ma believes that the groundbreaking discovery of unpaired Weyl

magnetic monopole can provide new insight into the fundamental physics of Weyl magnetic monopole in condensed matter. And the special properties of Weyl magnetic monopole, such as high electronic mobility, special response to an outside magnetic field and low heat rate, can make it a good candidate for spintronic computer devices like quantum computing and neuromorphic computing in the future.

More information: J.-Z. Ma et al, Observation of a singular Weyl point surrounded by charged nodal walls in PtGa, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-24289-0](https://doi.org/10.1038/s41467-021-24289-0)

Provided by City University of Hong Kong

Citation: Physicist observes the first unpaired Weyl magnetic monopole (2021, September 14) retrieved 4 May 2024 from <https://phys.org/news/2021-09-physicist-unpaired-weyl-magnetic-monopole.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.