

## Modeling improvements promise increased accuracy for epidemic forecasting

October 26 2021



For large R0 during the first wave, peak times are distributed around 55 days (most probable peak date) between 42 and 80 days. For the second wave at a lower R0, peak dates are distributed around 130 days on a much wider distribution as predicted by the study's theoretical and numerical model. Credit: Maxence Arutkin

Accurate forecasting of epidemic scenarios is critical to implementing effective public health intervention policies. While much progress has



been made in predicting the general magnitude and timing of epidemics, there's still room for improvement in forecasting peak times, as unfortunately evidenced with H1N1 and COVID-19, when peak times occurred later than predicted.

In *Chaos* researchers from France and Italy use dynamical stochastic modeling techniques to reveal that infection and <u>recovery</u> rate fluctuations play a critical role in determining peak times for epidemics.

"Some averaged quantities, like infection and recovery rates, are highly sensitive to parameter fluctuations, which means that the latter must be understood, even when average behavior is the only focus of interest," said co-author Maxence Arutkin. "Our work shows that <u>epidemic</u> peak timing depends on these fluctuations, and neglecting them in epidemiological models can lead to inaccurate epidemic scenarios and unsuitable mitigation policies, not to mention enable viruses to evolve into new variants."

Using a susceptible-infected-recovered epidemic model that incorporates daily fluctuations on control parameters, the study applies probability theory calculations to infection counts at the beginning of an epidemic wave and at peak times for populations in Italy. While previous works using standard epidemiological models have suggested there is a delay between the epidemic peak date and its prediction (without fluctuations), the researchers suggest the epidemic peak <u>time</u> depends not only on the mean value of the infection and recovery rates but also on their fluctuations.

To predict epidemic trajectory, an important parameter is the basic reproduction number, R0, which describes the average number of infections transmitted from an individual. Infection and recovery rate fluctuations lead to lognormal probability distribution of the number of infected people, similar in its analytical form to price distributions for



financial assets.

"In the short term, even when average infections transmitted from a single individual are less than one, we can observe epidemic resurgence due to parameter fluctuations," said Arutkin. "Also, a dispersion of the epidemic peak time can be quantified showing that, without taking these fluctuations into account, the peak time estimates are biased."

The study reveals that improved prediction depends on both R0 levels and fluctuations in <u>infection</u> and recovery rates and may provide policymakers with a tool to assess the consequences of parameter fluctuations based on different R0 levels.

"Our findings suggest we must introduce parameter fluctuations in epidemiological models going forward," said Arutkin.

**More information:** Maxence Arutkin et al, Delayed epidemic peak caused by infection and recovery rate fluctuations, *Chaos* (2021). doi.org/10.1063/5.0067625

## Provided by American Institute of Physics

Citation: Modeling improvements promise increased accuracy for epidemic forecasting (2021, October 26) retrieved 17 May 2024 from <u>https://phys.org/news/2021-10-accuracy-epidemic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.