


Crucial step identified in the conversion of biomass to methane

October 4 2021



Credit: DOI: 10.1073/pnas.2111682118

Microbial production of methane from organic material is an essential process in the global carbon cycle and an important source of renewable energy. This natural process is based on a cooperative interaction between different types of microorganisms: the fermenting bacteria and the methane-producing archaea. The former converts so-called primary fermentation products from biomass decomposition, including fatty acids into intermediate products such as acetic acid, formate or H₂.

Specialized archaea can then form methane from them. The syntrophic interaction of fermenting bacteria with methanogenic archaea is crucial

for the globally relevant conversion of biomass into methane. However, scientists have not yet been able to clarify how the oxidation of saturated [fatty acids](#) can be coupled with the thermodynamically extremely unfavorable reduction of CO₂ to methane and how such a process can enable the growth of both microorganisms involved. A research team from the University of Freiburg, the Technical University of Darmstadt and the University of Bern in Switzerland led by Prof. Dr. Matthias Boll from the Institute of Biology II at the University of Freiburg has now been able to uncover a crucial step in this process: they found the missing enzymatic link and its function, which makes methane formation from fatty acids traceable from an energetic standpoint. The researchers published their findings in the journal *Proceedings of the National Academy of Sciences*.

Study on oxidoreductase

The scientists investigated a previously uncharacterized membrane-bound oxidoreductase (EMO) from the fermenting bacterium *Syntrophus aciditrophicus*. They provided biochemical evidence that the heme-b cofactors of this membrane-bound oxidoreductase and a modified quinone with perfectly matched redox potentials are the main players in this microbial process. Bioinformatics analyses also suggest that these oxidoreductases are widely distributed in prokaryotes, organisms such as bacteria and [archaea](#) whose cells lack a nucleus. "The results not only close our knowledge gap on the conversion of biomass to [methane](#)," Boll explains. "We may additionally identify EMOs as previously overlooked key components of lipid metabolism in the vast majority of all microorganisms."

More information: Michael Agne et al, The missing enzymatic link in syntrophic methane formation from fatty acids, *Proceedings of the National Academy of Sciences* (2021). [DOI: 10.1073/pnas.2111682118](https://doi.org/10.1073/pnas.2111682118)

Provided by University of Freiburg

Citation: Crucial step identified in the conversion of biomass to methane (2021, October 4)
retrieved 3 May 2024 from

<https://phys.org/news/2021-10-crucial-conversion-biomass-methane.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.