

Researchers zero in on methane released from reservoirs

October 20 2021, by Sarah Derouin



Reservoirs like Lac des Dix, the reservoir of the Grande Dixence Dam in Hérémece, Switzerland, release large amounts of methane to the atmosphere. Credit: Jérémy Toma, CC-BY-SA-4.0

Methane is a potent greenhouse gas that accounts for about a fifth of today's global warming. In addition to methane emitted from livestock and other agricultural practices, the gas is also spewed from the production, transport, and burning of fossil fuels and the breakdown of

organic material. Methane is released into the atmosphere continually. Scientists track these emissions to use in climate modeling and greenhouse gas budgeting studies, so accurate measurements and emissions modeling systems are crucial.

Wetlands and inland water bodies like reservoirs contribute a substantial amount of [methane](#) to the atmosphere, but prior estimates are limited and highly variable. Prior reservoir emissions estimates ranged anywhere from 18 to 70 teragrams of methane per year.

Now a new study by Johnson et al. uses a grid approach to estimate daily methane emissions from reservoirs. Their approach gives detailed estimates of methane flux, both geographically and by time.

The authors note that the amount of methane released from reservoirs can vary depending on the area of the reservoir, local climate, time of day or year that flux measurements are taken, and freeze-thaw cycles. To better capture these fluctuations, the researchers set up a $0.25^\circ \times 0.25^\circ$ (latitude-longitude) [grid pattern](#) over reservoirs around the world. They used available methane measurements reported for reservoirs, correcting for differences created by taking measurements at a certain time of day or season. They used satellite observations to define ice-free emissions periods to model total global emissions.

Using the average methane flux estimates for each grid and the duration of emissions periods, the researchers calculated global daily emissions over a full yearly cycle. They found that the global surface area of reservoirs is nearly 300,000 square kilometers and that reservoirs are concentrated in temperate and tropical regions. Reservoirs in temperate and tropical regions had higher methane emissions (121.4–140.9 milligrams per square meter per day) compared with boreal reservoirs (39.1 milligrams per square meter per day), and warmer months produced higher emissions.

Globally, the researchers estimated that 10.1 teragrams of methane are emitted from reservoirs per year—lower than previous estimates. The authors note differences in surface areas, treatment of seasonal/daily emissions measurements, and shorter emissions periods to account for the differences. They note that their new methods help reduce the uncertainties in global methane budgets and that their estimates will be useful for bottom-up biogeochemical and top-down atmospheric methane emissions estimates.

More information: Matthew S. Johnson et al, Spatiotemporal Methane Emission From Global Reservoirs, *Journal of Geophysical Research: Biogeosciences* (2021). [DOI: 10.1029/2021JG006305](https://doi.org/10.1029/2021JG006305)

This story is republished courtesy of Eos, hosted by the American Geophysical Union. Read the original story [here](#).

Provided by American Geophysical Union

Citation: Researchers zero in on methane released from reservoirs (2021, October 20) retrieved 22 June 2024 from <https://phys.org/news/2021-10-methane-reservoirs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.