

Scientists pinpoint evolutionary genes that allow lizards to give birth like mammals

October 8 2021



Credit: Hans Recknagel

Scientists studying the evolution of birth in lizards, from egg-laying to live births, have pinpointed the evolutionary genes from which the species is evolving to 'build' a new mode of reproduction.

The study—led by the University of Glasgow and published in *Nature Ecology and Evolution*—found that a significantly similar amount of the same genes involved in the pregnancy of lizards were shared with other mammals and live-bearing vertebrates.

Evolving from egg-laying to live birth—also known as viviparity—is a major evolutionary step; however, it is almost impossible to study the genes that lead to such major changes because when animals have evolved live-birth it was usually in the distant past.

However, the so-called Common Lizard—a species found in much of Eurasia, including Scotland—only evolved to live-birth relatively recently.

All vertebrates are either egg-laying or live-bearing, yet the genes that determined being one or the other were not known.

For the study, the scientists spent several years studying rare hybrid offspring reproduced from breeding between egg-laying and viviparous (live birthing) lizards. The researchers tracked female reproduction in [natural populations](#) in remote regions in the Alps, then used advanced genome analysis to look at how and why these lizards' DNA differs.

The analyses revealed genetic profiles particular to live-bearing lizards included genes known to be associated with fertility and pregnancy in mammals. It also found that gene pathways of tissue growth and immune function used in [mammal](#) reproduction were active in the lizards. Genes in egg-laying traits included some involved in egg productivity in birds.

Dr. Hans Recknagel, who led the field and genome research during his Ph.D. and postdoctoral research, said: "This was fascinating research, not least because in this species of lizard egg-laying populations still occur and interbreed with live-bearing ones.

"Despite the 300 million years of evolutionary distance between reptiles and mammals, when we compared our findings across vertebrates—looking at seven independent origins of viviparity, or live birth, in mammals, squamates, and fish—we found that a significant

amount of the same genes are involved in pregnancy in lizards and in mammals."

Professor Kathryn Elmer, the study's senior author from the University of Glasgow's Institute of Biodiversity, Animal Health and Comparative Medicine, said: "Though the origins of viviparity, or live [birth](#), are independent the genetic similarities are unlikely to be a coincidence."

"This shows how [evolution](#) tends to find similar answers to the challenges organisms face. It can be using similar established genetic networks, as we found here, but also new and different [genes](#) can be involved, allowing some flexibility in how evolution 'builds' a new reproductive mode again and again."

The [study](#), "The functional genetic architecture of egg-laying and live-bearing reproduction in common [lizards](#)" is published in the journal *Nature Ecology and Evolution*. The work was primarily funded by the UK's Natural Environment Research Council.

More information: Hans Recknagel et al, The functional genetic architecture of egg-laying and live-bearing reproduction in common lizards, *Nature Ecology & Evolution* (2021). [DOI: 10.1038/s41559-021-01555-4](#)

Provided by University of Glasgow

Citation: Scientists pinpoint evolutionary genes that allow lizards to give birth like mammals (2021, October 8) retrieved 23 May 2024 from <https://phys.org/news/2021-10-scientists-evolutionary-genes-lizards-birth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.