

## Three-dimensionally oriented orbital angular momentum of light

October 21 2021



A wave packet contains the intersection of spatiotemporal and spatial vortices. The average orbital angular momentum per photon is accurately controllable in the value and orientation. Credit: Science China Press

Recently, *National Science Review* published online the research results on three-dimensionally oriented orbital angular momentum of light from Professor Qiwen Zhan's team at the University of Shanghai for Science and Technology.



Photonic angular <u>momentum</u> plays an important role in the interaction between light and matter. Photons can carry spin angular momentum related to the polarization and <u>orbital angular momentum</u> (OAM) related to the phase structure. Generally, the OAM direction is parallel to the propagation direction of light field, and the angular momentum carried by each photon is the product of the reduced Planck constant and topological charge.

Based on the collision between spatiotemporal vortices and spatial vortices, the researchers accomplished the accurate control of the threedimensional orientation of photonic OAM, and added a new degree of freedom to angular momentum of light. The average OAM carried by each photon in the spatiotemporal wave packet generated by the experiment is controllable not only in three-dimensional direction, but also in magnitude. The value of OAM per photon is not limited to the integer multiple of the reduced Planck constant, but can also be a non-integer multiple of the reduced Planck constant. Furthermore, a traveling wave packet can contain the collision of multiple spatiotemporal vortices and multiple spatial vortices, enabling fast modulation of the magnitude and direction of photonic OAM in the spatiotemporal domain.

This research achieves the accurate control of photonic OAM in threedimensional space and <u>time domain</u>, and thus is of great significance to the study of the interaction between <u>light</u> and metamaterials and the development of new photonic devices.





The intersection of spatiotemporal and spatial vortices is achieved based on the manipulation in the spatiotemporal frequency domain. Slice interference technique enables the reconstruction of three-dimensional optical field. Credit: Science China Press

**More information:** Chenhao Wan et al, Photonic orbital angular momentum with controllable orientation, *National Science Review* (2021). DOI: 10.1093/nsr/nwab149

Provided by Science China Press



Citation: Three-dimensionally oriented orbital angular momentum of light (2021, October 21) retrieved 8 May 2024 from <u>https://phys.org/news/2021-10-three-dimensionally-orbital-angular-momentum.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.