

Urbanites face heightened flood risk due to forest loss

October 21 2021



Credit: NASA

The devastating impact of flooding in Queensland's north, exacerbated by forest loss, is badly affecting urban areas according to University of Queensland-led research.



UQ's Dr. Jaramar Villarreal Rosas studied flooding in the Brigalow Belt Bioregion—just north of Townsville, to Emerald and Rockhampton—over a 13-year period.

"The capacity for <u>natural ecosystems</u> to mitigate flooding has significantly decreased across the Brigalow Belt due to deforestation," Dr. Rosas said.

"Forests help mitigate floods by reducing the amount of water flowing on the soil surface soon after a rain event.

"While overall flood mitigation continues to decrease, urban residents in towns like Townsville and Charters Towers have borne the brunt.

"Specifically, we found that, between 2002 and 2015, the decrease in <u>flood protection</u> to <u>urban residents</u> was double the amount we observed in any other sector, followed by rural communities and the food sector.

"This is a direct result of continued deforestation and land clearing of forests in the region, when forests are proven to reduce the intensity of flooding events."

The research team said the Brigalow Belt is home to animals and plants that exist nowhere else in the world.

Dr. Rosas said through conservation of natural areas and forests, nearby urban regions could be shielded from significant flooding.

"Through the protection of these forests we'll be able to reduce the intensity of floods, which cause billions of dollars of in damage to homes and businesses," she said.

"This study can be used to develop management strategies to address



these inequalities in the distribution of flood protection and other <u>ecosystem services</u>."

Moving forward, Dr. Rosas said the differences in how different people are affected by ecosystem degradation must be better reflected in policy.

"While restoration of some of these ecosystems can bolster flood protection, there must be special consideration when developing restoration plans to make sure inequalities are accounted for and addressed," she said.

"This means a special focus on restoring specific areas within the Brigalow Belt.

"In a further study, we're developing a planning approach that indicates the priority areas in this region, where applying restoration actions would deliver the greatest <u>flood protection</u> benefits to the most vulnerable population sectors, at the minimum cost."

Research senior author Professor Jonathan Rhodes said a greater focus on mitigating the impact of climate change is the best path forward.

"We have to place more emphasis on restoration and adaptation policies with a long-term focus," Professor Rhodes said.

"This includes reforestation and land clearing regulation, improved emergency response plans, and careful planning about where we locate infrastructure at risk of flooding."

The research has been published in Science of the Total Environment.

More information: Jaramar Villarreal-Rosas et al, The impacts of land use change on flood protection services among multiple beneficiaries,



Science of The Total Environment (2021). DOI: 10.1016/j.scitotenv.2021.150577

Provided by University of Queensland

Citation: Urbanites face heightened flood risk due to forest loss (2021, October 21) retrieved 28 June 2024 from <u>https://phys.org/news/2021-10-urbanites-heightened-due-forest-loss.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.