


A computer algorithm that speeds up experiments on plasma

November 16 2021, by Bob Yirka



Plasma ion temperature climbs rapidly after 1 ms and reaches values of ~1 keV by 3 ms into the discharge on Shot 46366. Typical temperatures are lower and show little to no rise in time. Credit: DOI: 10.1038/s41598-017-06645-7

A team of researchers from Tri Alpha Energy Inc. and Google has developed an algorithm that can be used to speed up experiments conducted with plasma. In their paper published in the journal *Scientific Reports*, the group describes how they plan to use the algorithm in nuclear fusion research.

As research into harnessing [nuclear fusion](#) has progressed, scientists have found that some of its characteristics are too complex to be solved in a reasonable amount of time using current technology. So they have increasingly turned to computers to help. More specifically, they want to adjust certain parameters in a device created to achieve fusion in a reasonable way. Such a device, most in the field agree, must involve the creation of a certain type of [plasma](#) that is not too hot or too cold, is stable, and has a certain desired density.

Finding the right parameters that meet these conditions has involved an incredible amount of trial and error. In this new effort, the researchers sought to reduce the workload by using a [computer](#) to reduce some of the needed trials. To that end, they have created what they call the "optometrist's [algorithm](#)." In its most basic sense, it works like an optometrist attempting to measure the visual ability of a patient by showing them images and asking if they are better or worse than other images. The idea is to use the crunching power of a computer with the intelligence of a human being—the computer generates the options and the human tells it whether a given option is better or worse.

The experiments by the researchers are being run on Tri Alpha Energy's C2-U computer, and thus far, the researchers report that the work is progressing much faster than it was before. One run, for example, now takes just eight minutes. Each run involves firing beams of hydrogen atoms at a sample of plasma. Doing so serves to keep the plasma

spinning inside of a [magnetic field](#) for as long as 10 milliseconds. The researchers are hoping that each run is bringing them closer to building a [fusion](#) reactor that generates more energy than it consumes.

More information: E. A. Baltz et al, Achievement of Sustained Net Plasma Heating in a Fusion Experiment with the Optometrist Algorithm, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-06645-7](https://doi.org/10.1038/s41598-017-06645-7)

© 2021 Science X Network

Citation: A computer algorithm that speeds up experiments on plasma (2021, November 16) retrieved 26 April 2024 from <https://phys.org/news/2021-11-algorithm-plasma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.