


Capturing the impact of human sewage on Earth's coastal ecosystems

November 10 2021



A) Global map of the terrestrial sources (green to blue) and coastal diffusion of inputs (yellow to purple) of total wastewater N, measured in $\text{log}_{10}(\text{gN})$ in both. Coastal plumes have been buffered to line segments to exaggerate patterns to be visible at the global scale. Insets show zoomed-in views of the B) Ganges, C) Danube, and D) Chang Jiang (Yangtze) Rivers, showing wastewater plumes at high resolution. Credit: Tuholske et al., 2021, PLOS ONE, CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)


A first-of-its-kind, high-resolution mapping analysis estimates the amounts of nitrogen and pathogens released into coastal ecosystems from human wastewater sources around the world. Cascade Tuholske (now affiliated with the Columbia Climate School) and colleagues at the University of California, Santa Barbara, present this research in the open-access journal *PLOS ONE* on November 10, 2021.

The researchers have created a [visual representation](#) of this, available [here](#).

Human sewage can introduce disease-causing pathogens and [nitrogen](#) into the ocean, potentially impacting human health as well as [coastal ecosystems](#) and the communities that depend on them for such purposes as fishing. However, most research into humans' impact on coastal ecosystems has focused on [agricultural runoff](#), while investigations on [human sewage](#) have been limited.

To better capture the impact of sewage on coastal ecosystems, Tuholske and colleagues conducted a novel analysis in which they estimated and mapped nitrogen and pathogen inputs into the ocean from sewage for about 135,000 watersheds around the world at a resolution of 1 kilometer. The assessment employed newly available, high-resolution data on global human populations and modeled how wastewater plumes entering the ocean would overlap with different ecosystems.

The analysis suggests that wastewater from human sewage introduces 6.2 teragrams of nitrogen into coastal ecosystems per year—for comparison, that is about 40 percent of estimated inputs from agriculture. Sixty-three percent of the nitrogen is from sewage systems, 5 percent from septic systems, and 32 percent from untreated, direct input.




The global total wastewater input is 6.2Tg N, with 3.9Tg from sewers, 0.3Tg from septic, and 2Tg from direct input. The top 40 countries are shown in the horizontal bar chart; remaining countries are in the pinwheel, grouped by continent or larger geographical region. Values for all countries are also reported in S5 Table in S1 File. Note that the Netherlands is shown in both places (in red) to help connect the scale of the two parts of the figure. Credit: Tuholske et al., 2021, *PLOS ONE*, CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)

Of the watersheds that appear to release the most nitrogen from sewage, most are located in India, Korea, and China, with the Chang Jiang (Yangtze) River contributing 11 percent of the global total. The researchers also identified hotspots for coral reef exposure to nitrogen in China, Kenya, Haiti, India, and Yemen. Seagrass exposure hotspots were found in Ghana, Kuwait, India, Nigeria, and China. The Chang Jiang and Brahmaputra Rivers have the highest input of pathogens.

Further research will be needed to refine the model and its estimates. Nonetheless, this work provides a new resource that could play a key role in efforts to mitigate harm to ecosystems and [human health](#)—such as by highlighting locations where tradeoffs between managing nitrogen and pathogen levels are particularly important to consider.


A




Coral



B

Seagrass



-  Habitat presence
-  No wastewater impact
-  High wastewater impact

Maps show where A) coral reefs and B) seagrass beds are heavily impacted (raster cells in top 2.5% of exposure; red dots), not impacted (no exposure to wastewater N; dark blue dots), or impacted but not in the top 2.5% (yellow dots). Raster cells are represented as points which visually over-represents the habitat; red is overlaid on top which makes it visually dominant; blue points are transparent and overlaid on green/yellow points such that higher densities of unimpacted areas are brighter blue. Credit: Tuholske et al., 2021, *PLOS ONE*, CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)

The authors add: "The sheer scale of how much wastewater is impacting coastal [ecosystems](#) worldwide is staggering. But because we map wastewater inputs to the ocean across more than 130,000 watersheds, our results identify target priority areas to help marine conservation groups and public health officials to work together and reduce the impacts of wastewater on coastal waters across the planet."

More information: Tuholske C, Halpern BS, Blasco G, Villasenor JC, Frazier M, Caylor K (2021) Mapping global inputs and impacts from of human sewage in coastal ecosystems. *PLoS ONE* 16(11): e0258898. [journals.plos.org/plosone/arti ... journal.pone.0258898](https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0258898)

Provided by Public Library of Science

Citation: Capturing the impact of human sewage on Earth's coastal ecosystems (2021, November 10) retrieved 30 January 2023 from <https://phys.org/news/2021-11-capturing-impact-human-sewage-earth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.