

Researcher argues for existence of companion candidate stars to the eclipsing binary Algol

November 4 2021, by Riitta-Leena Inki



(a) Algol's O-C data (red circles). The time-axis units are days between November 1782 and October 2018. The green continuous line shows the five signal model for the first 226 years of data before the dotted vertical line. The data minus model differences are offset from zero to -0.3 (blue circles). (b) The last 15 years of O-C data. Beyond the vertical dotted line begins the prediction for the last 10 years (continuous green line). The dotted green lines show the prediction error limits. This test checks how well does the model for the first 226



years data predict the last 10 years of data. The prediction is excellent. Credit: University of Helsinki

Dr. Lauri Jetsu from the University of Helsinki has analyzed observations of Algol. He argues that Algol has many companion stars which have not been detected from earlier observations. The results have been published in *The Astrophysical Journal*.

Algol is an eclipsing binary, where the two <u>stars</u> Algol A and Algol B orbit around their common center of mass. Their <u>orbital period</u> is 2.867 days. The abbreviation for this binary system is Algol AB.

Algol's primary eclipses occur when the dimmer Algol B partially covers the brighter Algol A. These primary eclipses last ten hours, and they can be observed with naked eye. Goodricke (1783) determined Algol's orbital period from naked eye observations of these events. The primary eclipses would be repeated regularly exactly after 2.867 days, if nothing disturbed the motions of Algol AB binary system. All these future eclipses could be computed from the multiples of the constant period 2.867 days.

The presence of a third member Algol C in this multiple star system was confirmed in late 1950s. Algol C and Algol AB orbit around their common center of mass. One round takes 1.86 years. The orbital motions of Algol C and Algol AB cause a light travel time effect. We observe the primary eclipses earlier when Algol AB is closer to us, and later when Algol AB is further away from us. During every 1.86 years round, Algol C causes the same regular positive and negative time shifts in the observed eclipse epochs of Algol AB. The range of these time shifts is only about nine minutes. Due to these time shifts, the observed eclipse epochs (O=Observed) differ from the computed constant period



eclipse epochs (C=Computed). These differences are called the O-C data.

There may be even four or five new companions

Lauri Jetsu analyzed the O-C data of Algol between November 1782 and October 2018. He applied his recently formulated discrete chi-square method to these data. This method is designed for detecting regular periodic signals. These detections succeed even if the signals are superimposed on an irregular aperiodic trend. From O-C data of Algol AB, the discrete chi-square method can detect the light travel time effect signals of five or six companion star candidates. The O-C data alone can not be used to establish the exact number of these candidates. One of these candidates is the known "old" companion Algol C. The orbital periods of the other four or five "new" companion star candidates are between 20 and 219 years.

"These stars are candidates until new observations confirm their existence," says Jetsu. He also shows that the periodic signals of these candidates can predict the observed Algol's O-C changes.





Lauri Jetsu has studied Algol for a long time and tattooed in his hand a hieroglyph which describes Algol. Ancient Egyptian called Algol as Horus, which was a manifestation from the god and from the king. Credit: Riitta-Leena Inki

Why have these candidates not been detected earlier?

Algol is so close to the sun that astronomers can observe its eclipses with naked eye. Algol's new companion candidate stars would be literally in our backyard.

"The paradox is that Algol is 'too bright," says Jetsu. Algol can hide these new companion candidate stars even from our most powerful modern space telescopes, just like our sun can hide all other stars during daytime, says Jetsu. He points out that, for example, the cutting-edge



equipment onboard the Gaia satellite could not detect Algol's new companion candidates. Jetsu argues that future interferometric observations may be used to directly confirm the existence of at least some of these new Algol's companion candidates.

More information: Lauri Jetsu, Say Hello to Algol's New Companion Candidates, *The Astrophysical Journal* (2021). <u>DOI:</u> <u>10.3847/1538-4357/ac1351</u>

Provided by University of Helsinki

Citation: Researcher argues for existence of companion candidate stars to the eclipsing binary Algol (2021, November 4) retrieved 6 May 2024 from <u>https://phys.org/news/2021-11-companion-candidate-stars-eclipsing-binary.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.