

Fate of sinking tectonic plates revealed

November 11 2021



Researchers including The University of Texas at Austin have published a study in *Nature* that says subducting tectonic plates become segmented like a slinky snake, in a process similar to geologic boudinage (pictured) but on a much larger scale. Credit: Haakon Fossen, Structural Geology/Cambridge University Press

Our world's surface is a jumble of jostling tectonic plates, with new ones emerging as others are pulled under. The ongoing cycle keeps our

continents in motion and drives life on Earth. But what happens when a plate disappears into the planet's interior?


The question has long puzzled scientists because conventional wisdom said that sinking tectonic plates must remain intact to keep pulling on the portion behind it, but according to geophysical evidence, they are destroyed.

Now, in a study published Nov. 11 in *Nature*, scientists say they've found an answer that reconciles the two stories: Plates are significantly weakened as they sink but not so much that they break apart entirely.

The finding came after scientists put tectonic plates through a computer-generated gauntlet of destructive geologic forces. The [model](#) showed that as the plate enters the mantle, it bends abruptly downward, cracking its cold, brittle back. At the same time, the bending changes the fine grain structure of the rock along its underbelly, leaving it weakened. Combined, the stresses pinch the plate along its weak points, leaving it mostly intact but segmented like a slinky snake.

This means the plate continues to be pulled under despite becoming folded and distorted.

According to the researchers, the model predicted a scenario that matches observations from Japan. Studies of the region where the Pacific tectonic plate dives—or subducts—under Japan have turned up large cracks where the plate bends downward, and they have shown signs of weaker material underneath. Deep seismic imaging conducted by The University of Texas at Austin's Steve Grand has also revealed tectonic shapes in the Earth's mantle under Japan that appear a close match for the slinky snake in the model.



The researchers put sinking tectonic plates through a simulated gauntlet of

destructive geologic forces. Only when all forces were included did the plate behave in a way that matches geophysical evidence from the surface. Credit: Taras Gerya, David Bercovici, Thorsten Becker/Springer Nature.

Co-author Thorsten Becker, a professor in UT's Jackson School of Geosciences, said that the study does not necessarily close the book on what happens to subducting plates, but it certainly gives a compelling case to explain several important geologic processes.

"It's an example of the power of computational geosciences," said Becker who assisted in developing the model and is a faculty associate at UT's Oden Institute for Computational Engineering & Sciences. "We combined these two processes that geology and rock mechanics are telling us are happening, and we learned something about the general physics of how the Earth works that we wouldn't have expected. As a physicist, I find that exciting."

The study's lead author, Taras Gerya, a professor of geophysics at ETH Zurich, added that until now, geophysicists had lacked a comprehensive explanation for how [tectonic plates](#) bend without breaking.

Things got interesting when the researchers ran their simulations with a hotter interior, similar to the early Earth. In these simulations, the tectonic snake segments made it only a few miles into the mantle before breaking off. That means that subduction would have occurred intermittently, raising the possibility that modern [plate tectonics](#) began only within the past billion years.

"Personally, I think there are a lot of good arguments for plate tectonics being much older," Becker said, "but the mechanism revealed by our model suggests things might be more sensitive to the temperature of the

mantle than we thought, and that, I think, could lead to interesting new avenues of discussion."

Becker and Gerya were joined by David Bercovici, a geophysicist at Yale University whose investigation into how rock grains are altered in the [deep mantle](#) helped motivate the research. The study is based on a two-dimensional computer model of plate tectonics incorporating Bercovici's rock deformation research and other [plate](#)-weakening mechanics. The researchers are now studying the phenomena using 3D models and plan to investigate what those models can tell them about the occurrence of earthquakes.

More information: T. V. Gerya et al, Dynamic slab segmentation due to brittle–ductile damage in the outer rise, *Nature* (2021). [DOI: 10.1038/s41586-021-03937-x](#)

Provided by University of Texas at Austin

Citation: Fate of sinking tectonic plates revealed (2021, November 11) retrieved 26 April 2024 from <https://phys.org/news/2021-11-fate-tectonic-plates-revealed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.