

What's in a flame? The surprising mystery of how soot forms

November 17 2021, by Andrew Breeson



Credit: Unsplash/CC0 Public Domain

Soot is one of the world's worst contributors to climate change. Its impact is similar to global methane emissions and is second only to carbon dioxide in its destructive potential. This is because soot particles absorb solar radiation, which heats the surrounding atmosphere, resulting in warmer global temperatures. Soot also causes several other environmental and health problems including making us more

susceptible to respiratory viruses.

Soot only persists in the atmosphere for a few weeks, suggesting that if these emissions could be stopped then the air could rapidly clear. This has recently been demonstrated during recent lockdowns, with some major cities reporting clear skies after industrial emissions stopped.

But [soot](#) is also part of our future. Soot can be converted into the useful carbon black product through thermal treatment to remove any harmful components. Carbon blacks are critical ingredients in batteries, tires and paint. If these carbons are made small enough they can even be made to fluoresce and have been used for tagging [biological molecules](#), in catalysts and even in solar cells.

Given the importance of soot and how long humankind has been producing it, you would think its formation was completely understood. However, this is not the case. In particular, the critical transition when the [molecules](#) cluster to form the very first nanoparticles of soot is unknown.


If the origins of soot were to be entirely understood, we could potentially eliminate its formation and therefore drastically reduce its [environmental impact](#) as well as make better carbon materials. With this in mind, researchers from the University of Cambridge and [Cambridge CARES](#) have recently published a comprehensive review on the birth of soot—where molecules become particles.

In the review, titled: "Soot inception: Carbonaceous nanoparticle formation in flames" published in [Progress in Energy and Combustion Science](#), the authors [Dr. Jacob Martin](#), [Dr. Maurin Salamanca](#) and CARES Director [Professor Markus Kraft](#) begin by noting that;

"It has only been in the last decade, however, that experimental and

computational techniques in combustion science have been able to peek behind the door to reveal insights into the earliest formation mechanisms of carbonaceous particulates in the flame."


The figure below shows some of these new experimental insights along the path from fuel to soot. In this diagram it is nanoparticle formation (soot inception) that is the birth of the soot particle.



The graphical abstract from "Soot inception: Carbonaceous nanoparticle formation in flames" Credit: Jacob Martin

Two main pathways have been suggested for soot inception—either physical condensation in which molecules form droplets or chemical

polymerisation in which molecules react to form particles. But either pathway by itself is non-optimal, as "physical and electrical condensation of precursor molecules is rapid but too weak to hold soot together, while most chemical bonds are strong but the mechanisms proposed to date are too slow to account for rapid growth of soot as observed in experiments."



Schematic of various soot nanoparticles arranged as a function of their C/H ratio and molecular weight. Credit: Jacob Martin

Instead, the authors suggest a "middle way" involving mechanisms with both physical and chemical aspects. Promising options are highlighted involving π -radicals and diradicals, however, conclusive evidence for a specific mechanism as well as predictive models are still lacking.

Ultimately, the authors conclude that "the emission of carbonaceous nanoparticles needs to be a research and industrial priority for the future of combustion devices and new material applications."

"Soot inception: Carbonaceous nanoparticle formation in flames" is published in *Progress in Energy and Combustion Science* by researchers from Cambridge Centre for Advanced Research and Education in Singapore Ltd and University of Cambridge.

More information: Jacob W. Martin et al, Soot inception: Carbonaceous nanoparticle formation in flames, *Progress in Energy and Combustion Science* (2021). [DOI: 10.1016/j.pecs.2021.100956](https://doi.org/10.1016/j.pecs.2021.100956)

Provided by Cambridge Centre for Advanced Research and Education in Singapore

Citation: What's in a flame? The surprising mystery of how soot forms (2021, November 17) retrieved 8 May 2024 from <https://phys.org/news/2021-11-flame-mystery-soot.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.