

Using neutrons to analyse human remains

November 1 2021



Figure 1. INS spectra (measured in MAPS, with 5240 cm–1 incident energy) of human femur burned at temperatures between 200 and 650°C, either under aerobic (a) or anaerobic (b) conditions. The temperature range corresponding to the largest spectral variations is highlighted by a dashed rectangle. (The spectra of reference calcium hydroxyapatite (HAp, SRM 2910b) and of femur unburned (unb) and burned at 700°C [28] are also shown, for comparison purposes.). Credit: DOI: 10.1098/rsos.210774

A UK neutron facility has been used to develop a technique to help



better understand human skeletal remains that have been subject to heating.

Just as we do today, our ancient ancestors practiced cremation in their funeral rites, which means that <u>skeletal remains</u> are often found in a burned state.

However, determining whether the <u>bone</u>'s heating is the result of cremation practices or even ritualistic cannibalism is difficult to determine.

Researchers have used the Science and Technology Facilities Council's ISIS Neutron and Muon Source to develop a new technique for archeological or forensic investigators to tackle this problem.

Bouncing neutrons

The team, led by the University of Coimbra, Portugal, used the ISIS facility to bounce neutrons off samples of human bone subjected to varying degrees of heating.

The study aimed to analyze <u>human bones</u> exposed to temperatures between 200°C and 650°C.

At this low end of the heating scale, the experiment recreates the sort of temperatures achieved in cooking vessels, which provides information about potential cannibalism or de-fleshing rituals.

Physical and chemical changes

Studying this temperature range is important because the physical and <u>chemical changes</u> that can occur when bone is buried can actually mimic



the effects of low-intensity heating.

This makes it difficult to differentiate between buried and moderatelyheated bone.

Unfortunately studying bone exposed to temperatures in this range is challenging.

The porosity of bone means that those heated to around 500 °C are more fragile than those exposed to higher temperatures.

Electrically neutral neutrons

This is where ISIS's <u>neutron</u> beams come in.

Neutrons are used in this sort of research because they are electrically neutral, which means they can pass into and through materials without causing damage.

Dr. Stewart Parker, senior research scientist at ISIS, said: "This was a fascinating piece of research to be involved in and really demonstrates the breadth of applications of neutron science. Neutrons provide a non-destructive way of gaining unique insights into materials from catalysts and materials for the electronics industry to biological samples. The collaboration between ISIS and the University of Coimbra in Portugal goes back over 20 years. The current project has allowed us to get detailed information on how, and why, bones change through burning, both helping us understand our ancestry and explore potential forensic applications."

Impact of oxygen



The data obtained by the team showed that structural changes within the bone varied depending on the amount of oxygen to which the sample was exposed.

The greater the oxygen exposure, the lower temperature at which heatinduced changes, such as fat and protein destruction, occurred.

The latest study builds on the team's previous work that looked at the effects of more extreme heating on the structure of human bone.

The team now has data that spans a <u>temperature</u> range of 200°C to 1000°C.

Comprehensive catalog

All this provides a comprehensive catalog of heat-induced changes.

It will potentially allow archaeologists to determine whether human remains have been exposed to lower temperatures during cooking, or to higher temperatures during cremation.

It is also possible that the research could be used to assist in the interpretation of the bone mineralisation processes that occur during fossilization.

This may help to better understand the biology of extinct vertebrates.

Forensic applications

Along with the archeological applications, this work could also be applied to forensic investigations where bone is found that has been subjected to fire.



When human remains are found at a crime scene, or the site of an accident, the first priority of forensic investigators is to figure out the identity of the person they belong to.

When this can't be done using superficial characteristics, such as facial structure and hair color (what they look like), forensic pathologists will resort to:

- dental records
- fingerprints
- DNA.

However, should the remains consist of the skeleton alone, it may not be possible to determine the identity so easily and the priority shifts to identifying:

- sex
- age at time of death
- general physical characteristics, such as height.

Measuring bone dimensions

This is done by measuring the dimensions of the bones and crossreferencing those dimensions with known markers for age and sex, such as the width of pelvic bones.

It might then be possible to match these parameters with those of missing persons.

This becomes much more difficult, however, if the remains had been subjected to intense heat, as they might in a plane crash or house fire.



Heated bones warp and shrink

When bones are heated they warp and shrink, which changes their dimensions, making the age, sex and size much more difficult to determine.

This work will allow scientists to gauge exactly what changes occur in the bone's structure when it subjected to heat.

The idea being that, if you know how the bone changes as it burns, you will be able to "turn the clock back" and extrapolate a bone's original unburned shape from its burned shape.

More information: M. P. M. Marques et al, The impact of moderate heating on human bones: an infrared and neutron spectroscopy study, *Royal Society Open Science* (2021). <u>DOI: 10.1098/rsos.210774</u>

Provided by Medical Research Council

Citation: Using neutrons to analyse human remains (2021, November 1) retrieved 17 July 2024 from <u>https://phys.org/news/2021-11-neutrons-analyse-human.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.