


Pseudobulbs: sink for exogenous nitrogen in epiphytic orchids

November 5 2021, by Li Yuan



Development of *Pleione aurita* over a whole growing season. a, Dormant pseudobulb; b, flowering stage; c, full leaf expansion; and d, leaf shedding. Credit: ZHANG Wei

Pseudobulbs, which are somewhat enlarged or thickened succulent stems, are common in epiphytic orchids. Nitrogen can be transported from old to young pseudobulbs and between leaves and pseudobulbs.

Quantifying [nitrogen](#) allocation and resorption of pseudobulbs is important for understanding the survival and adaptation strategies of orchids.

Researchers from Xishuangbanna Tropical Botanical Garden (XTBG) and Kunming Institute of Botany (KIB) of the Chinese Academy of Sciences quantified the nitrogen allocation and resorption in the epiphytic orchid *Pleione aurita* in relation to [nitrogen supply](#). They found that both the old and newly produced pseudobulbs can act as a sink for exogenous nitrogen, but with much lower sink activity in the former.

Their study was published in *Scientia Horticulturae*.

The researchers investigated the physiological performance, biomass allocation and nitrogen resorption of *P. aurita*, an ornamental and medicinal orchid plant, under three nitrogen levels. Using ^{15}N labeling, they compared the contributions of exogenous nitrogen and stored nitrogen, and the allocation to each organ.

They found that the leaf took up the largest proportion of both fertilizer and old pseudobulb-stored nitrogen. The fruit preferred to use pseudobulb-stored nitrogen regardless of exogenous nitrogen supply. The resorption of nitrogen in the leaf and pseudobulb was more efficient than other plant organs, but decreased with increasing nitrogen availability.

"Our results suggested an important role of the pseudobulb in the nitrogen economy of *P. aurita*, and the high dependence of the

reproduction process on the pseudobulb-stored nitrogen rather than an exogenous supply," said Fan Zexin, a corresponding author of the study.

More information: Wei Zhang et al, Quantifying the nitrogen allocation and resorption for an orchid pseudobulb in relation to nitrogen supply, *Scientia Horticulturae* (2021). [DOI: 10.1016/j.scienta.2021.110580](https://doi.org/10.1016/j.scienta.2021.110580)

Provided by Chinese Academy of Sciences

Citation: Pseudobulbs: sink for exogenous nitrogen in epiphytic orchids (2021, November 5) retrieved 11 May 2024 from <https://phys.org/news/2021-11-pseudobulbs-exogenous-nitrogen-epiphytic-orchids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.