

# How to turn specific genes on and off

November 9 2021

---



Credit: CC0 Public Domain

All the cells in an individual's body bear the same genetic code. It is the reading and writing of this code—the "turning on" and "turning off" of specific genes in specific cells—that gives the cells their identities. Imagine, for example, the disastrous situation wherein the genes that encode the digestive enzymes of the stomach would be turned on in the retinal cells of the eye and begin to devour the surrounding tissue. One of the ways by which a cell turns off specific genes is by the reversible addition to the DNA of a minuscule chemical called a methyl group at the precise location of that specific gene.

Scientists know that genes with more of this "DNA methylation" tend to be "turned off" and that genes with less of this methylation tend to be "turned on." But because until now it hasn't been possible to manipulate the levels of DNA methylation at [specific genes](#), there remain many questions about what specific instances of DNA methylation do, how they contribute to normal cellular function, and how their dysregulation contributes to disease.

In a recent study, published in *Nature Communications*, McGill University researchers demonstrate how they have managed to remove specific instances of DNA methylation at specific genes in mouse and [human cells](#) grown in culture by using CRISPR/Cas9 genome editing technology. They show that this DNA "de-methylation" activity can be targeted to anywhere in the DNA—any gene of interest—without editing the [genetic code](#), and with no off-target activity at undesired locations in the DNA. The researchers also describe the approaches needed to produce complete removal of DNA methyl marks in the hopes that scientists around the world can use this new technique to start to discover actionable instances where genes that should be on have been turned off by DNA methylation—such as, for example, the insulin gene in diabetes—and use this technique to establish new paradigms for disease

treatment.

**More information:** Daniel M. Sapozhnikov et al, Unraveling the functional role of DNA demethylation at specific promoters by targeted steric blockage of DNA methyltransferase with CRISPR/dCas9, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-25991-9](https://doi.org/10.1038/s41467-021-25991-9)

Provided by McGill University

Citation: How to turn specific genes on and off (2021, November 9) retrieved 24 June 2024 from <https://phys.org/news/2021-11-specific-genes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.