


Catalyst technology converts methane greenhouse gas into useful, valuable chemicals

December 8 2021



Credit: Pixabay/CC0 Public Domain

Methane, which produces more warming than other greenhouse gasses and is the subject of newly announced U.S. emission restrictions, is hard to break down and keep out of the atmosphere.

It's not that the primary component of natural gas is chemically complex. A methane molecule is just one carbon atom and four hydrogens.

But those carbon-hydrogen bonds are hard to break.

Typically, that involves [high temperatures](#) and mixing the flammable gas with oxygen to produce syngas to make methanol and hydrogen to make ammonia. That's expensive and potentially explosive.

Other conversion reactions aren't very efficient and also produce the most abundant of the [greenhouse gasses](#), carbon dioxide.

So, is there another way to use methane and keep it out of the atmosphere? One that's safe and efficient? One that returns value? One that helps fight climate change?

Yue Wu and his research group at Iowa State University have been looking for ways to do all that over several years of experimental failures and discoveries. Now, they and a group of collaborators have found and tested a [catalyst technology](#) that appears to be an answer.

"The results provided a potential solution to this long-time challenge and represented the best stability, conversion rate, and selectivity to convert methane to ethane or ethylene, two main precursors for the modern petrochemical industry," according to a project summary written by Wu, the Herbert L. Stiles Professor in Chemical Engineering at Iowa State.

A research paper recently published in the journal *Nature Catalysis* reports the discovery. Wu and Yang Xiao, a senior research engineer in the Davidson School of Chemical Engineering at Purdue University in Indiana, are corresponding authors. Zhe Li, who earned a doctoral degree at Iowa State in 2019 and is now a postdoctoral fellow at Johns Hopkins University in Maryland, is the first author. (See sidebar for

additional co-authors.)

The Iowa State University Office of Innovation Commercialization is seeking a patent for the technology.

Failure leads to success

The catalyst consists of one or two layers of platinum, each layer just an atom thick, deposited on two-dimensional metal carbide structures called "MXenes." In this case, the structures are made from carbon, molybdenum and titanium.

Wu said his research group discovered the thin layers essentially allow every platinum atom to be used as a catalyst and prevents the formation of residues that cover and deactivate the platinum. That means less platinum is required to make the catalyst.

Wu said his group started studying carbides—combinations of carbon and metals—about five years ago with support from the Office of Naval Research. The original work was to identify the electrical and thermal properties of various carbides. But the work didn't go as expected—the material's thermal conductivity was much lower than predicted.

"You can think of this as a failure," Wu said.

But, the researchers discovered the MXene surfaces are very active and able to absorb many molecules. And so, with support from Wu's Stiles professorship and the Iowa State College of Engineering, Wu's research group began studying these MXenes as a potential catalyst.

"We had never seen carbide so active," Wu said. "It's usually very inert. It's used, for example, for high-speed drill bits—the surface is hard and inert."

They started using the technology to remove hydrogen from shale gas, the natural gas found in shale rock formations. That work evolved to study other reactions involving natural gas.

"Nobody tried to use these carbides for these high-volume reactions before," Wu said.

Keys to the methane-to-ethane/ethylene conversion are making the carbides pure enough and making the surfaces clean enough to support the reactions, Wu said. Get it all right, and those reactions exhibit about 7% methane conversion with about 95% selectivity toward ethane/ethylene in a continuously operating fixed-bed reactor. The products can be turned into plastics and resins, such as the common and ubiquitous polyethylene plastic.

"Remarkably, these novel catalysts run for 72 hours of continuous operation without any signs of deactivation, indicating a promising start toward technologies suitable for exploitation on the industrial scale," Wu wrote.

That's all very good news, he said.

Methane emissions, after all, are such a contributor to climate change that world leaders took steps to restrict them during COP26, the recent United Nations summit on climate change in Glasgow, Scotland. President Joe Biden announced the U.S. will take steps to restrict methane emissions from existing oil and gas operations. More than 100 countries also signed a Global Methane Pledge to reduce methane emissions by 30% over the next nine years.

The researchers' new catalyst technology could advance those efforts to keep methane out of the atmosphere. Wu calls the technology "revolutionary," saying it "opens the door to reducing the emission of

[methane](#) and its combustion product, CO₂, in the future."

More information: Zhe Li et al, Direct methane activation by atomically thin platinum nanolayers on two-dimensional metal carbides, *Nature Catalysis* (2021). [DOI: 10.1038/s41929-021-00686-y](https://doi.org/10.1038/s41929-021-00686-y)

Provided by Iowa State University

Citation: Catalyst technology converts methane greenhouse gas into useful, valuable chemicals (2021, December 8) retrieved 23 May 2024 from <https://phys.org/news/2021-12-catalyst-technology-methane-greenhouse-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.