

Researchers find climate change record in clam shells

December 14 2021



Archaeologists have discovered a way to reconstruct past climates by analyzing the shells of this species of short-lived surf clams, Donax obesulus, which have been consumed by people since ancient times. Credit: Jacob Warner, LSU

The tiny, pale surf clam about the size of a fingernail that most people have seen and collected on beaches around the world holds clues in its



shell to Earth's past. For the first time, researchers have been able to identify the monthly, and even weekly, ocean temperatures recorded in these smooth clam shells. Because ancient civilizations consumed these ubiquitous clams and left the shells at archaeological sites, researchers now have a new way to reconstruct climate and its fluctuations from nearly 3,000 years ago.

"The equipment available now, compared to the past, is precise and powerful enough to be able to reveal the sea surface temperature and the overall climate at a specific location when the clam was building its shell. This gives us archaeologists and paleoclimatologists another tool in our proverbial toolbox to reconstruct past climate. As we know today, climate can influence all kinds of practices and behaviors, which may have been the case in <u>ancient civilizations</u> as well," said LSU Department of Geography & Anthropology doctoral candidate Jacob Warner, who is the lead author on this new research published in *Chemical Geology*.

Warner will present this new research, 'Trace Element Ratios (Sr/Ca, Mg/Ca, and Ba/Ca) in the Short-Lived Bivalve Donax obesulus: Potential Environmental Proxies?' at the American Geophysical Union, or AGU, Fall meeting on Tuesday, Dec. 14 from 4:00–6:00 p.m. Central. Members of the media can register to attend AGU for free.

Warner and colleagues are focused on tracking a climate phenomenon that affects a large part of the world called the El Niño Southern Oscillation, or ENSO, the warm phase of which is characterized by warmer than normal <u>ocean temperatures</u>, increased rainfall and more tropical storms and hurricanes in the southern U.S.





LSU Ph.D. candidate Jacob Warner drilled along the clam's shell to collect samples at each interval of time during the clam's lifespan to get a snapshot of the ocean temperature as the shell grew. Credit: Jacob Warner, LSU

Warner's study sites are in northern Peru, which is one of the areas most impacted by ENSO. The cold, deep water off the coast of Peru makes it one of the most productive fisheries in the world; however, when a warm ENSO event occurs, it can result in fishery die-offs.

To capture the <u>sea surface temperature</u> from different phases of ENSO, the researchers collected 18 surf clams from markets and coastal beaches in 2012, 2014 and 2016. This new research used a species of short-lived surf clam called Donax obesulus, which has not been used to reconstruct climate before. Previous studies have successfully used the short-lived intertidal clam species, Mesodesma donacium. However, this species is now extinct in northern Peru.

Like <u>trees</u> and <u>tree rings</u>, clams create layers in their shells as they grow. Warner drilled along the shell to collect samples at each interval of time during the clam's lifespan to get a snapshot of the ocean temperature as the shell grew.





Archaeological site, Caylán, in the Nepeña Valley, Peru. Credit: Jacob Warner, LSU

In a separate, forthcoming paper, Warner collaborated with fellow archeologist Aleksa Alaica, a postdoctoral researcher at the University of Alberta to analyze the surf <u>clam</u> Donax obesulus found at an archaeological site in the Jequetepeque Valley in northern Peru. They discovered that the clams' shells were larger during warmer ENSO events; therefore, shell size can also be used as a paleoclimate proxy. They also discovered that the ancient people, who lived at this site, preferentially harvested larger individual clams, which indicates a



fisheries management practice in place more than 2,000 years ago.

"This collaborative study showcases the power of simple methods to examine complex questions. Our work highlights the multiple factors affecting <u>shell</u> growth and the resilience of communities in the past to respond to climate and social transformations," Alaica said.

More information: Jacob P. Warner et al, Investigating the influence of temperature and seawater δ 18O on Donax obesulus (Reeve, 1854) shell δ 18O, *Chemical Geology* (2021). DOI: 10.1016/j.chemgeo.2021.120638

Provided by Louisiana State University

Citation: Researchers find climate change record in clam shells (2021, December 14) retrieved 26 June 2024 from <u>https://phys.org/news/2021-12-climate-clam-shells.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.