

New crystal structure for hydrogen compounds for high-temperature superconductivity

December 13 2021



Figure 1. Clathrate structures of $LaYH_{12}$ and LaY_3H_{24} . Using computer simulations, a global team of researchers have predicted new crystal structures for realizing a high-temperature superconductor with ternary hydrides, a step toward realizing a low-cost and lossless power transmission technology. Credit: Ryo Maezono from JAIST.

Superconductivity is the disappearance of electrical resistance in certain



materials below a certain temperature, known as "transition temperature." The phenomenon has tremendous implications for revolutionizing technology as know it, enabling low-loss power transmission and maintenance of electromagnetic force without electrical supply. However, superconductivity usually requires extremely low temperatures ~ 30 K (the temperature of liquid nitrogen, in comparison, is 77 K) and, therefore, expensive cooling technology. To have a shot at realizing a low-cost superconducting technology, superconductivity must be achieved at much higher transition temperatures.

Materials scientists have had a breakthrough on this front with crystalline materials containing hydrogen, known as "metal hydrides." These are compounds formed by a metal atom bonded with hydrogen that have been predicted and realized as suitable candidates for achieving even room-temperature superconductivity. However, they require extremely high pressures to do so, limiting their practical applications.

In a new study published in *Chemistry of Materials*, a group of researchers led by Professor Ryo Maezono from Japan Advanced Institute of Science and Technology (JAIST) performed <u>computer</u> <u>simulations</u> to expand the search for high-temperature superconductors, looking for <u>potential candidates</u> among ternary hydrides (hydrogen combined with two other elements).

"In ternary hydrides, the number of elements is increased from two to three. While this enormously increases the number of possible combinations and can make the problem of predicting suitable materials more difficult, it also increases our chances of coming across a potential high-temperature superconductor," explains Prof. Maezono.

Using the supercomputer at the university, the researchers examined



possible crystal structures for (LaH6) (YH6)y compounds (y=1-4), looking for configurations that would yield stable structures, allowing their synthesis in the laboratory at high pressures. Starting from a random structure, the simulations went through various possible combinations of elements, testing their stability at extremely high pressures ~ 300 GPa.

The simulations revealed clathrate (Cmmm-) structures of $LaYH_{12}$ and LaY_3H_{24} , consisting of LaH_{24} and YH_{24} cages stacked on top of each other (Figure 1), as viable candidates for high-temperature and high-pressure superconductors. "The longer stacking for Cmmm-LaY_3H_{24} lead to a slightly increased transition temperature," explains Prof. Maezono. Among the possible structures, the highest transition temperature (145.31 K–137.11 K) was observed for LaY_3H_{24} . The researchers attributed the origin of higher transition temperature to a high "density of states" and high "phonon frequency," two parameters that are used to assess superconductivity in materials.

These findings have excited the researchers, who optimistically speculate the discovery of more such <u>high-temperature superconductors</u>. "It is quite possible to predict using simulations other new combinations of elements that would improve the desired properties further," says Prof. Maezono.

More information: Peng Song et al, High-Tc Superconducting Hydrides Formed by LaH24 and YH24 Cage Structures as Basic Blocks, *Chemistry of Materials* (2021). DOI: 10.1021/acs.chemmater.1c02371

Provided by Japan Advanced Institute of Science and Technology

Citation: New crystal structure for hydrogen compounds for high-temperature superconductivity



(2021, December 13) retrieved 6 May 2024 from <u>https://phys.org/news/2021-12-crystal-hydrogen-compounds-high-temperature-superconductivity.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.