

How a floating fern withstands rain

December 21 2021



A drop of water on a leaf of the tropical floating fern *Salvinia molesta*. Credit: University of Tübingen

The tropical floating fern *Salvinia molesta* has developed sophisticated structures to allow water to roll off its leaves quickly—even during

heavy rainfall. This relieves the pressure on the leaves floating on the water surface, but even more importantly, it keeps the stomata open for air exchange. This allows the fern to absorb carbon dioxide which is essential for photosynthesis. This was discovered in an interdisciplinary collaborative project between the working group of Professor James Nebelsick and Dr. Dr. Wilfried Konrad from the Department of Geosciences at the University of Tübingen and Dr. Anita Roth-Nebelsick from the State Museum of Natural History in Stuttgart. In their experiments, the team simulated rainfall and used high-speed cameras to track the path of water droplets slapping onto the fern's leaves. The leaf structures are efficiently adapted to rainfall and may contribute to the strong growth and rapid spread of the fern. Outside its native Brazil, *Salvinia molesta* is considered a dangerous invasive plant, displacing other species. The new study is published in the *Journal of the Royal Society Interface*.

Salvinia molesta, like all floating ferns, floats freely on the water surface. Of each trifoliate whorl, two floating leaves lie on top, with the plant extending one root-like leaf downward. This [fern](#) quickly forms thick, dense mats on the [water surface](#). It originates from humid tropical regions in Brazil, where it is regularly exposed to [heavy rainfall](#). The upper surface of the floating leaves is covered with highly water-repellent hairs, whose structure resembles tiny egg-whisks. "These trichomes ensure that the submerged leaves do not become waterlogged," reports Wilfried Konrad. "Other aquatic plants, however, achieve this with much less elaborate structures. We therefore asked ourselves what benefit the fern obtains from the little whisks."

Experiments simulate rainfall

The research team obtained specimens of *Salvinia molesta* from the Tübingen Botanical Gardens and the Wilhelma in Stuttgart. In one experiment, leaves of the plant were fixed in place while drops of water

of varying size splashed onto them from different heights. "Overall, the leaves quickly shook off most of the artificial rain from the surface. The camera images show that the trichomes respond elastically to the kinetic energy of the falling drops," Konrad says. Even more important in shaking off the water, however, is the elasticity of the leaves as a whole. The rain experiment was also carried out on free-floating leaves, underlining the key role of elasticity. "When the leaves were fixed, it took much longer for their surface to become water-free again," Konrad explains.

The researchers observed how [water droplets](#) bounced off the leaves, with some taking on a pan-cake-like shape or being broken up into smaller droplets. "Especially in the experiments with large water droplets falling from a great height onto the leaves, we noticed that there was always some residual water left under the tiny whisks," says Konrad. "Surprisingly, the trichomes seemed to rather prevent the water from beading off quickly."

Water-repellent structures

Detailed observations, however, revealed that they were very efficient at keeping the leaf base free of water. "The energy state of a droplet is more favorable on the trichomes than between them. This is due to the water-repelling effect of the whisks," Konrad explains. When a drop of water rolls over the trichomes of the leaf, it literally sucks up the residual water trapped between the trichomes due to its strong surface forces. "This keeps water off the stomata at the base of the floating fern's leaves, which are also covered with water-repellent nanocrystals made of wax. This is where the gas exchange that is vital for the plant takes place." Konrad added that this may be a decisive factor in the high productivity of the *Salvinia molesta*, which can double its leaf mass within a few days under favorable conditions.



Experimental setup for the drop height of water drops of 430 centimeters in the stairwell in the paleontological collection (Sigwartstr. 10, Tübing-en). At the upper end of the tube the water drops were sent on their journey, at the bottom near the floating fern leaves the high-speed camera is set up. Credit: University of Tübingen

More information: Wilfried Konrad et al, The impact of raindrops on *Salvinia molesta* leaves: effects of trichomes and elasticity, *Journal of The Royal Society Interface* (2021). [DOI: 10.1098/rsif.2021.0676](https://doi.org/10.1098/rsif.2021.0676)

Provided by University of Tübingen

Citation: How a floating fern withstands rain (2021, December 21) retrieved 26 June 2024 from <https://phys.org/news/2021-12-fern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.