

# Too dry, too hot, or too wet: Increasing weather persistence in European summer

December 6 2021

---



Credit: CC0 Public Domain

"In our study, we show that persistent weather conditions have an increasing similarity in summer over the North Atlantic, Europe and Siberia, favoring more pronounced extreme weather events. In Europe alone, about 70% of the land area is already affected by more persistent

weather situations," says Peter Hoffmann from the Potsdam-Institute for Climate Impact Research (PIK), lead author of the study published in *Nature's Scientific Reports*. "This means that people, especially in densely populated Europe, will likely experience more and also stronger and more dangerous weather events."

## **Prolonged sunny or rainy conditions lead to extreme events**

To prove this, the scientists analyzed the persistence of specific [weather](#) conditions. They applied established image comparison methods on atmospheric data, comparing millions of successive weather circulation patterns worldwide over the last 40 years. They especially looked at two individual extreme events, the 2010 [heat wave](#) over Russia and the extraordinary dry summer over Europe in 2018. "We found that weather patterns in general are more persistent now than some decades ago," says Hoffmann. "Especially in summer, heat waves often last longer now, and also rainfall events tend to linger longer and to be more intense. The longer these weather conditions last, the more intense the extremes can become, both on the warm and dry side as well as on the steady rain side."

The rise in persistent weather conditions is to a large extent due to dynamical changes in the atmosphere as the [westerly winds](#) tend to stop pushing forward weather systems which therefore become more persistent, turning some sunny days into heatwaves of several weeks as well as intensive rainfall into floods.

## **Climate models might underestimate rise in persistent weather patterns**

Like the eye of a knowledgeable observer, the new image comparison

method systematically scans atmospheric fields and helps to assess to which extent successive weather circulation patterns change over time or remain stable. Fred Hattermann, co-author of the study and also scientist at PIK, explains the potential of the new approach: "Our method crucially improves the interpretation of long-term climate impacts. When applying the same methodology to climate model output, we do not see a comparable increase in weather persistence, especially not over Europe. Climate models may have been a bit too conservative, underestimating the rise in weather persistence—thus underestimating weather extremes over Europe." This becomes ever more important with ongoing climate change.

"Recent events and ongoing research have shown that the number and intensity of extreme events have increased due to the global rise in temperature," concludes Hattermann. "To counteract this worrying trend, the goal has to be to reduce [global greenhouse gas emissions](#) and by doing so limit climate change."

**More information:** Peter Hoffmann et al, Atmosphere similarity patterns in boreal summer show an increase of persistent weather conditions connected to hydro-climatic risks, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-01808-z](https://doi.org/10.1038/s41598-021-01808-z)

Provided by Potsdam Institute for Climate Impact Research

Citation: Too dry, too hot, or too wet: Increasing weather persistence in European summer (2021, December 6) retrieved 13 March 2024 from <https://phys.org/news/2021-12-hot-weather-persistence-european-summer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---