

Plasma lensing discovered in black widow pulsar





The upper panel shows the total intensity of pulse emission vs. pulsar spin and orbital phases with the sub-integration of 1 s of PSR J1720-0533. Enlargements of the ingress and egress of the pulsar are shown in the middle left and middle right panels, respectively. The bottom panels show the pulse flux density variations near the eclipse. Credit: XAO

Using the Five-hundred-meter Aperture Spherical radio Telescope (FAST), a research team led by Dr. Wang Shuangqiang from the Xinjiang Astronomical Observatory (XAO) of the Chinese Academy of Sciences discovered plasma lensing phenomenon in a black widow pulsar



PSR J1720-0533.

Black widow pulsar systems have a low-mass companion star in a compact orbit with a <u>millisecond pulsar</u>. They are characterized by ablating the companion by emission from <u>pulsar</u>. Black widow pulsars get their name from the "black widow" spiders, the females of which eat the males after mating. Black widow pulsars offer valuable opportunities to investigate the characteristics of the companion stars under intense irradiation.

In this study, the researchers found that the emission of PSR J1720-0533 during the ingress of the eclipse shows quasi-periodic modulations, which may be caused by plasma lensing.

By analyzing the lensing phenomenon, the researchers concluded that the maximum magnification for the lens is 1.6, corresponding to a lens size of tens of kilometers. The discovery of the plasma lensing phenomenon in PSR J1720-0533 demonstrates a link between the dispersion measurement and lensing.

Moreover, the researchers examined the polarization profiles near the eclipse of PSR J1720-0533 and found that the linear polarization of the emission disappeared before the dispersion measurement showed significant changes. This <u>phenomenon</u> provides strong evidence that there is a significant magnetic field in the companion.

These results suggest that magnetic fields play an important role in the eclipsing mechanism of black widow pulsars.

In addition, the researchers estimated the mass-loss rate of the companion to be 10^{-12} M \odot /yr and speculated that the companion will be destroyed completely in 10^{10} yr.



More information: S. Q. Wang et al, Unusual Emission Variations Near the Eclipse of Black Widow Pulsar PSR J1720–0533, *The Astrophysical Journal Letters* (2021). DOI: 10.3847/2041-8213/ac365c

Provided by Chinese Academy of Sciences

Citation: Plasma lensing discovered in black widow pulsar (2021, December 30) retrieved 26 June 2024 from <u>https://phys.org/news/2021-12-plasma-lensing-black-widow-pulsar.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.