

# Plastic-degrading enzymes increasing in correlation with pollution

December 14 2021

---



Credit: Unsplash/CC0 Public Domain

The number of microbial enzymes with the ability to degrade plastic is growing, in correlation with local levels of plastic pollution. That is the finding of a new study from Chalmers University of Technology,

Sweden, that measured samples of environmental DNA from around the globe. The results illustrate the impact plastic pollution is having on the environment, and hint at potential new solutions for managing the problem.

The problems of global plastic pollution are all too widespread, as mass-production of plastic has exploded in the last 70 years or so—from around 2 million tons per year to around 380 million. This has given sufficient evolutionary time for various microbes present in the environment to respond to these compounds, and many different enzymes have been discovered in previous studies with the ability to degrade different plastics.

The new study, recently published in the scientific journal *mBIO*, analyzed samples of environmental DNA from hundreds of locations around the world. The researchers used computer modeling to search for [microbial enzymes](#) with plastic-degrading potential, which was then cross-referenced with the official numbers for plastic waste pollution across countries and oceans.

"Using our models, we found multiple lines of evidence supporting the fact that the global microbiome's plastic-degrading potential correlates strongly with measurements of environmental plastic pollution—a significant demonstration of how the environment is responding to the pressures we are placing on it," says Aleksej Zelezniak, associate professor in systems biology at Chalmers University of Technology.

## **More enzymes in the most polluted areas**

In other words, the quantity and diversity of plastic-degrading enzymes is increasing, in direct response to local levels of [plastic pollution](#). In total, over 30,000 enzyme homologues were found with the potential to degrade 10 different types of commonly used plastic. Homologues are

members of protein sequences sharing similar properties. Some of the locations that contained the highest amounts were notoriously highly polluted areas, for example samples from the Mediterranean Sea and South Pacific Ocean.

"Currently, very little is known about these plastic-degrading enzymes, and we did not expect to find such a large number of them across so many different microbes and environmental habitats. This is a surprising discovery that really illustrates the scale of the issue," explains Jan Zrimec, first author of the study and former post-doc in Aleksej Zelezniak's group, now a researcher at the National Institute of Biology in Slovenia.

## **Potential value for fighting the global plastic crisis**

Every year around 8 million tons of plastic escapes into the world's oceans. The natural progresses for plastic degradation are very slow—the lifetime of a PET-bottle, for example, can be up to hundreds of years. The growth and accumulation of plastic waste in the oceans and on land is a truly global problem and there is an increasing need for solutions to manage this waste. The researchers believe that their results could potentially be used to discover and adapt enzymes for novel recycling processes.

"The next step would be to test the most promising enzyme candidates in the lab to closely investigate their properties and the rate of plastic degradation they can achieve. From there you could engineer microbial communities with targeted degrading functions for specific polymer types," explains Aleksej Zelezniak.

The article, *Plastic-Degrading Potential across the Global Microbiome Correlates with Recent Pollution Trends*, is published in the journal *mBio*. It was written by Jan Zrimec, Mariia Kokina, Sara Jonasson,

Francisco Zorrilla and Aleksek Zelezniak.

## **More evidence of a connection from comparing land and sea**

- The plastic-degrading enzymes which the researchers found were widely distributed across both [ocean](#) and soil microbiomes—a microbiome is the entire set of microbes present in a specific environmental location or group of locations.
- A lot of variation was encountered in the number and type of plastic particles, as well as plastic-degrading enzymes, found between the land and ocean samples. For example, the land samples contained many more phthalate-based plastic additive compounds. These are commonly used in all sorts of processes, and are known to be particularly susceptible to leaking during production, disposal, and recycling—processes which take place mainly on land. In correlation, more enzymes which were able to degrade these plastics were indeed found in the land samples, indicating a connection between the two.
- The ocean samples also revealed how the quantity of enzymes with degrading capability increased with depth. The dataset involved 67 locations from eight oceans, at three different depths, and showed consistent higher levels of degrading enzymes at deeper levels—again, indicating a connection to the greater level of microplastics that have been repeatedly observed at deeper levels in the ocean.

## **More information about how the researchers cataloged the enzymes**

- The researchers compiled a data set of 95 previously known enzymes with plastic-degrading or modifying capability. They

then used 'Hidden Markov Models' to search through data taken from some of the largest global metagenomic studies to identify homologous sequences from 236 locations. The researchers used samples of the internal human microbiome as a control for false positives—no plastic-degrading enzymes have as yet been identified within humans, despite concerns of microplastic-ingestion.

- They identified a total of around 30,000 enzyme hits, circa 12,000 in the ocean microbiome and 18,000 in the soil, corresponding to 10 major commercial plastics, including six polymers and four additives. Nearly 60% of the identified plastic-degrading enzymes did not map to any known [enzyme](#) classes, suggesting that the researchers uncovered novel [plastic](#)-degrading functional content.

**More information:** Jan Zrimec et al, Plastic-Degrading Potential across the Global Microbiome Correlates with Recent Pollution Trends, *mBio* (2021). [DOI: 10.1128/mBio.02155-21](https://doi.org/10.1128/mBio.02155-21)

Provided by Chalmers University of Technology

Citation: Plastic-degrading enzymes increasing in correlation with pollution (2021, December 14) retrieved 12 May 2024 from <https://phys.org/news/2021-12-plastic-degrading-enzymes-pollution.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.