

# **Soils in old-growth treetops can store more carbon than soils under our feet**

December 14 2021

---



Organic soil formed from epiphyte mats on a tree branch in a tropical humid lower montane rainforest at the Texas A&M University Soltis Center for Research and Education, Costa Rica. Credit: H. Connuck

New research reveals a previously underappreciated way old-growth forests have been recycling and storing carbon: Treetop soils. Branches in forest canopies can hold caches of soil that may store substantially more carbon than soils on the ground beneath them, and scientists are just beginning to understand how much carbon canopy soils—which exist on every continent except Antarctica—could store.

The new research on these unique soils, being presented on Wednesday, 15 December at 5:00 p.m. CST at AGU Fall Meeting 2021, marks the first attempt to quantify carbon capture by canopy soils. The work highlights another way [old-growth forests](#) are rich, complex ecosystems that cannot be quickly replaced by replanting forests.

Tree branches collect fallen tree leaves and other organic material over hundreds of years, like the ground does. On top of the [branches](#), the plant litter decomposes as it accumulates, forming a carbon-rich layer that can be several inches thick. The researchers climbed up into the rainforest canopy in Costa Rica, instruments in hand, to find out just how much carbon canopy soils can contain.

Active carbon, a short-term storage pool of organic carbon, was three times higher in canopy soil compared to soils underfoot, the researchers found.

"We knew these would be really organic-rich soils, but we didn't expect the extremely large amount of carbon compared to mineral soils," said Hannah Connuck, an undergraduate researcher at Franklin and Marshall

College who will be presenting the study results.




At 30 m height above the ground, Hannah Connuck collects canopy soil for carbon measurements from a humid lower montane rainforest. Credit: K. Munoz Elizondo

The researchers are still calculating the total concentration of organic carbon at their research site, but other research has found canopy soils to have up to 10 times higher concentrations of organic carbon, according to soil scientist Peyton Smith, a study co-author and Connuck's [soil](#) science mentor at Texas A&M University.

Connuck and Smith also measured how much carbon dioxide was being released by microbial organisms living in the canopy soils, which is critical for knowing whether soils are storing or releasing carbon overall. They found that even though the [microbes](#) were releasing higher volumes of carbon dioxide than ground soils, their rate of carbon storage

was rapid enough to compensate, likely making canopy soils a net carbon sink that has not been considered in carbon models yet.

"It could be a substantial [carbon](#) sink, and we need to account for it," Smith said.



Peyton Smith, lead researcher, climbs high to study canopy soils at the Texas A&M University Soltis Center for Research and Education, Costa Rica. Credit: H. Connuck

Like other soils, canopy soils take a long time to form, and therefore take a long time for a forest to recover if an area of old-growth is cut down. The soils also host unique microbiomes, including highly diverse microbial organisms and canopy-specific plants like epiphytic orchids.

"It's a good argument for keeping primary and other old-growth forests around, rather than harvesting and replanting with secondary growth

forests," Connuck said.

**More information:** Hannah Connuck et al, Canopy Soil Capture: An Overlooked Source or Sink of Carbon in Humid Tropical Forests? (2021). Available at [agu.confex.com/agu/fm21/meeting/paper.cgi/Paper/1001199](https://agu.confex.com/agu/fm21/meeting/paper.cgi/Paper/1001199)

Provided by American Geophysical Union

Citation: Soils in old-growth treetops can store more carbon than soils under our feet (2021, December 14) retrieved 10 May 2024 from <https://phys.org/news/2021-12-soils-old-growth-treetops-carbon-feet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.