


Team develops kilohertz label-free non-contact quantitative mapping of optical properties for strongly turbid media

December 27 2021



a, Spatially modulated light patterns are generated by the DMD and projected

onto the sample. Remitted light is collected by a camera. b, A Monte Carlo based model is used to extract optical absorption (μ_a) and reduced scattering (μ_s') from calibrated diffuse reflectance (R_d) at two spatial frequencies, in this case 0 mm⁻¹ and 0.1 mm⁻¹. c, Raw reflectance images are demodulated and calibrated to extract the spatial frequency response of the sample (i.e., diffuse reflectance). The Monte Carlo based inversion model is used to extract optical properties (absorption and reduced scattering) on a pixel-by-pixel basis. Here, an 8 × 6 cm region of human hand was imaged at 650 nm and the extracted optical absorption and reduced scattering maps are shown for the measured wavelength. d, Conventional 8-bit continuous-tone illumination pattern is converted to halftone 1-bit pattern which increases the maximum projection rate from 290 Hz to 23 kHz, by approximately two orders of magnitude. e, Raw images are collected for the same turbid sample region (i.e., human hand) with the conventional continuous-tone and the proposed halftone illumination patterns, respectively. While the binary discontinuity is visible in the halftone patterns after zoomed-in, the corresponding collected images have identical appearance with the ones from continuous-tone patterns as a result of low-pass filtering effect of turbid media. Credit: Yanyu Zhao, Bowen Song, Ming Wang, Yang Zhao, and Yubo Fan

The ability to quantify optical properties (i.e., absorption and scattering) of strongly turbid media has major impacts on the characterization of biological tissues, fluid fields, and many others. However, the task of quantitative imaging of optical properties for strongly turbid media is intrinsically challenging, as photon scattering hinders direct measurement over length scales larger than the mean free path. Few current imaging technologies can quantify absorption and scattering properties of strongly turbid media in a wide-field non-contact format, and none have been shown to map quantitative optical properties with high-speed (e.g., kHz) capabilities.

In a new paper published in *Light: Science & Application*, Professor Yanyu Zhao, Professor Yubo Fan, and co-workers at Beihang University

developed a new optical imaging method that for the first time can quantify [optical properties](#) of strongly turbid [media](#) as well as functional chromophore concentrations in tissue in a multi-kHz high-speed, label-free, non-contact, and wide-field manner. Their work was based on an emerging diffuse optical technology named spatial frequency domain imaging (SFDI). It can quantify optical properties of turbid media in a label-free, non-contact manner, and its camera-based detection scheme makes it intrinsically wide-field. It projects spatially modulated sinusoidal light patterns of different phases onto the sample using a digital-micromirror-device (DMD) and collects the reflectance images.

However, current SFDI technologies utilize a continuous-tone strategy and generate those sinusoidal patterns with 8-bit grayscale, which correspondingly has a limited maximum projection speed of 290 Hz determined by the DMD hardware. Consequently, while SFDI typically requires 5 projection patterns for the measurement of optical properties at a single wavelength, current SFDI technologies are severely limited for high-speed applications.

To address the bottleneck of measurement speed, the researchers proposed a halftone strategy to significantly increase the speed of SFDI by approximately two orders of magnitude, with no extra cost or modification on system hardware. Specifically, they first demonstrated the sinusoidal patterns generated by the halftone strategy for 1-bit DMD projection, which led to a maximum projection rate of 23 kHz, approximately two orders of magnitude faster than that in current SFDI technologies. They then validated the proposed halftone-SFDI against conventional SFDI measurements through experiments on an array of optical phantoms with a wide range of optical properties as well as in vivo human tissue. They also demonstrated dynamic monitoring of wide-field optical properties and functional chromophore concentrations in the rat brain cortex with the proposed halftone-SFDI. Enabled by the proposed method, they finally demonstrated kHz high-speed dual-

wavelength monitoring of wide-field optical properties of a highly-dynamic flow field.

"To the best of our knowledge, this is the first demonstration of label-free non-contact imaging of quantitative optical properties of strongly turbid media with a speed of kilohertz," they noted.

"This work has several important implications for scientific research and engineering applications. For example, mapping brain cortex functions is of fundamental significance for brain science, neuroscience, and cognitive psychology. With shortwave-infrared wavelengths, the halftone-SFDI can provide kHz wide-field quantification of water content in the turbid combustion flow, which could have a substantial impact on the design and optimization of those engines and dramatically reduce related energy cost." they added.

"The presented technique can spatially map quantitative optical absorption and scattering properties with a maximum speed of 7.6 kHz in strongly turbid media, and can map absolute functional chromophore concentrations in tissue with a maximum speed of 3.8 kHz. This breakthrough could open a new venue for both fundamental research and translational applications including brain science and fluid dynamics." the scientists forecast.

More information: Yanyu Zhao et al, Halftone spatial frequency domain imaging enables kilohertz high-speed label-free non-contact quantitative mapping of optical properties for strongly turbid media, *Light: Science & Applications* (2021). [DOI: 10.1038/s41377-021-00681-9](https://doi.org/10.1038/s41377-021-00681-9)

Provided by Chinese Academy of Sciences

Citation: Team develops kilohertz label-free non-contact quantitative mapping of optical properties for strongly turbid media (2021, December 27) retrieved 25 June 2024 from <https://phys.org/news/2021-12-team-kilohertz-label-free-non-contact-quantitative.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.