


Black holes and dark matter—are they one and the same?

January 4 2022, by Jim Shelton



A new study theorizes that primordial black holes formed after the Big Bang (the far left panel) constitute all dark matter in the universe. At early epochs they cluster and seed the formation of early galaxies and then eventually grow by feeding off gas and merging with other black holes to create the supermassive black holes seen at the center of galaxies like our own Milky Way today. Credit: Yale and ESA

Primordial black holes created in the first instants after the Big

Bang—tiny ones smaller than the head of a pin and supermassive ones covering billions of miles—may account for all of the dark matter in the universe.

That's the implication of a new model of the early [universe](#) created by astrophysicists at Yale, the University of Miami, and the European Space Agency (ESA). If proven true with data from the soon-to-launch James Webb Space Telescope, the discovery would transform scientists' understanding of the origins and nature of both dark matter and black holes.

Dark matter—which has never been directly observed—is thought to constitute the majority of matter in the universe and act as the unseen scaffolding upon which galaxies form and develop. Physicists have spent years testing a variety of dark matter candidates, including hypothetical particles such as sterile neutrinos, Weakly Interacting Massive Particles (WIMPs), and axions.

Black holes, on the other hand, have been observed. A black hole is a point in space where matter is so tightly compacted it creates intense gravity. Not even light can resist its pull. Black holes are found at the centers of most galaxies.

The new study, accepted for publication in *The Astrophysical Journal*, harkens back to a theory first proposed in the 1970s by physicists Stephen Hawking and Bernard Carr. At the time, Hawking and Carr argued that in the first fraction of a second after the Big Bang, tiny fluctuations in the density of the universe may have created an undulating landscape with "lumpy" regions that had extra mass. These lumpy areas would collapse into black holes.

Although the theory did not gain traction within the wider scientific community—the new study suggests that, if modified slightly, it could

have explanatory power after all.

If most of the [primordial black holes](#) were "born" at a size roughly 1.4 times the mass of Earth's sun, they could potentially account for all dark matter, said Yale professor of astronomy and physics Priyamvada Natarajan, the paper's theorist.

Natarajan and her colleagues say their new model shows that the first stars and galaxies would have formed around black holes in the early universe. Also, she said, primordial black holes would have had the ability to grow into [supermassive black holes](#) by feasting on gas and stars in their vicinity, or by merging with other black holes.

"Primordial black holes, if they do exist, could well be the seeds from which all supermassive black holes form, including the one at the center of the Milky Way," Natarajan said.

"What I find personally super exciting about this idea is how it elegantly unifies the two really challenging problems that I work on—that of probing the nature of dark matter and the formation and growth of black holes—and resolves them in one fell swoop," she added.

The James Webb telescope's mission will be to find the first galaxies that formed in the early universe and see stars forming planetary systems.

The new study's first author is Nico Cappelluti, a former Yale Center for Astronomy & Astrophysics Prize postdoctoral fellow who is now an assistant professor of physics at the University of Miami. Günther Hasinger, ESA's director of science, is the study's second author.

"Our study shows that without introducing new particles or new physics, we can solve mysteries of modern cosmology from the nature of dark [matter](#) itself to the origin of super-massive black holes," Cappelluti said.

Primordial black holes also may resolve another cosmological puzzle: the excess of infra-red radiation, synced with X-ray radiation, that has been detected from distant, dim sources scattered around the universe.

Natarajan and her colleagues said growing, primordial black holes would present "exactly" the same radiation signature.

Best of all, the existence of primordial black holes can be proven—or disproven—in the near future, courtesy of the James Webb Space Telescope and ESA's Laser Interferometer Space Antenna (LISA) mission announced for the 2030s.

If [dark matter](#) is comprised of primordial black holes, more stars and galaxies would have formed around them in the early universe—precisely the epoch that the James Webb telescope will be able to see. LISA, meanwhile, will be able to pick up gravitational wave signals from early mergers of primordial [black holes](#).

"If the [first stars](#) and galaxies already formed in the so-called 'dark ages,' Webb should be able to see evidence of them," Hasinger said.

Natarajan added, "It was irresistible to explore this idea deeply, knowing it had the potential to be validated fairly soon."

More information: Exploring the high-redshift PBH- Λ CDM Universe: early black hole seeding, the first stars and cosmic radiation backgrounds, arXiv:2109.08701 [astro-ph.CO]
arxiv.org/abs/2109.08701

Provided by Yale University

Citation: Black holes and dark matter—are they one and the same? (2022, January 4) retrieved 2

May 2024 from <https://phys.org/news/2022-01-black-holes-dark-matterare.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.