

How dairy farmers can adapt to climate change

January 5 2022, by Jeff Mulhollem



Dairy farms—especially in the Northeast—are increasingly subject to more stringent regulations to reduce nutrient losses. With expected warmer conditions that will result in increased ammonia volatilization from manure and more frequent and more severe storms that will cause more soluble phosphorus runoff, new strategies such as manure injection (shown) are needed to limit nitrogen and phosphorus losses from crop fields. Credit: Robert Meinen

Dairy farmers in the Northeast—facing a warming climate that exacerbates nutrient pollution but lengthens the growing season—can reduce the environmental impact of their operations and maximize revenues by double cropping and injecting manure into the soil, rather than broadcasting it.

That's the conclusion of a team of researchers, led by Penn State agroecologists, whose new study evaluated whole-farm production and the environmental and economic impacts of adopting these practices on a representative [dairy](#) farm in central Pennsylvania under recent historical and projected mid-century climate.

The research is important, according to Heather Karsten, associate professor of crop production/ecology in the College of Agricultural Sciences, because [dairy farms](#)—especially in the Northeast—are increasingly subject to more stringent regulations to reduce nutrient losses. With expected warmer conditions that will result in increased ammonia volatilization from manure and more frequent and more severe storms that will cause more soluble phosphorus runoff, new strategies are needed to limit nitrogen and phosphorus losses from crop fields.

To reach their conclusions, researchers simulated farm management strategies using the Integrated Farm System Model developed by C. Alan Rotz, an agricultural engineer with the U.S. Department of Agriculture's Agricultural Research Service. The strategies were informed by a long-running, dairy cropping systems experiment at Penn State's Russell E. Larson Agricultural Research Center, supported by the USDA. That simulation enabled the scientists to determine the effects on crop yields, feed production, nitrogen losses, sediment erosion, sediment-bound and soluble losses of phosphorus, greenhouse gas emissions, fossil energy use, and production costs.




The difference between a field treated with injected liquid manure (left) and broadcasted manure is obvious in this comparison photo. Injecting manure reduces ammonia volatilization and soluble phosphorus runoff from crop fields. Credit: Robert Meinen, Penn State

In findings recently published in *Agricultural Systems*, the researchers reported that double cropping and subsurface injection of manure reduced total [nitrogen](#) losses by 12%-18% and total phosphorus losses by 16%-19%. Adoption of these strategies, they noted, provided a feasible adaptation and mitigation approach for future climate by reducing projected increases in soluble [phosphorus](#) runoff and ammonia

emissions caused by warmer temperatures and greater precipitation.

Significantly, these benefits can be achieved while maintaining and potentially reducing total farm-production costs, Karsten pointed out.



Environmental benefits of double cropping and manure injection can be achieved while maintaining and potentially reducing total farm-production costs. Credit: Heather Karsten, Penn State

"There's a lot of interest in double cropping among Northeast dairy [farmers](#)," she said. "I think farmers are realizing they have a longer

[growing season](#), and there are more precipitation events in spring that make getting into the field to do any operation more challenging. Our results demonstrate that adoption of these strategies can help mitigate dairy farm environmental impacts now and even more in the future."

José Castaño-Sánchez, former postdoctoral researcher in the Department of Plant Science at Penn State, currently a postdoctoral research associate at New Mexico State University and the USDA, and Rotz contributed to the research.

More information: José P. Castaño-Sánchez et al, Double cropping and manure management mitigate the environmental impact of a dairy farm under present and future climate, *Agricultural Systems* (2021). [DOI: 10.1016/j.agsy.2021.103326](#)

Provided by Pennsylvania State University

Citation: How dairy farmers can adapt to climate change (2022, January 5) retrieved 15 June 2024 from <https://phys.org/news/2022-01-dairy-farmers-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.