


Study inspects the origin of high-mass X-ray binary 4U 2206+54

January 31 2022, by Tomasz Nowakowski



The 3D trajectories of 4U 2206+54/BD+53 2790 and BD +53 2820 \equiv Gaia EDR3 2005418950349782272, a member of Cep OB1 association, in

Galactocentric Cartesian coordinates in the past. Credit: Hambaryan et al., 2022.

European astronomers have observed a peculiar high-mass X-ray binary known as 4U 2206+54. Results of the research provide essential information regarding the origin of this system. The study was presented in a paper published January 19 on the arXiv pre-print repository.

X-ray binaries are composed of a normal star or a white dwarf transferring mass onto a compact neutron star or a black hole. Based on the mass of the companion star, astronomers divide them into low-mass X-ray binaries (LMXBs) and high-mass X-ray binaries (HMXBs).

4U 2206+54 is an HMXB hosting a neutron star accreting from the wind of its companion, designated BD +53 2790, which exhibits a radial velocity modulation. Its [orbital period](#) of just 9.5 days is one of the shortest among the known HMXB systems.

With the strength of the surface characteristic magnetic field at a level of at least 20 trillion G, BD +53 2790 is assumed to be a magnetar—a rare, strongly magnetized neutron star. The donor star in this binary is a peculiar O9 V star with an overabundance of helium.

However, the origin of 4U 2206+54 remains unknown. Thus, in order to shed more light on this matter, a team of astronomers led by Valeri Hambaryan of the Astrophysical Institute and University Observatory Jena in Germany, has conducted a kinematic study of this HMXB.

"In order to identify the possible birthplace of 4U 2206+54 one needs to determine its possible membership to a stellar group either currently or in the past. The latter also requires to perform their trace back motion study in the Galaxy to test the concept: 4U 2206+54 and a stellar group

or some of its members in the past were 'in the same place at the same time,'" the researchers wrote in the paper.

The study found that HMXB 4U 2206+54/BD+53 2790 and a star known as BD+53 2820 that is the brightest star of one of the subgroups of the Cepheus OB1 (Cep OB1) association were at the at the same place (about 11,200 light years away) some 2.8 million years ago. The researchers suppose that at this location and time, a supernova (SN) in a close massive binary took place and can be considered as the birthplace of the currently observed binary.

Furthermore, the studied HMXB is currently at a distance of approximately 10,100 [light years](#) and has a space velocity of 75–100 km/s with respect to BD+53 2820. These parameters suggest that the progenitor of the neutron star hosted by 4U 2206+54 lost some 4 to 9 solar masses during the supernova explosion and the [neutron](#) star of this binary received a kick velocity of about 200–350 km/s. The researchers estimate that the progenitor's initial mass was at a level of at least 32 solar masses.

"Our analysis of motion shows that 4U 2206+54 originates in the OB association Cep OB1, from which it escaped about 2.8 ± 0.4 Myr ago due to the SN of 4U 2206+54's progenitor," the authors of the paper concluded.

More information: V. Hambaryan et al, The origin of the high-mass X-ray binary 4U 2206+54/BD +53 2790. arXiv:2201.07770v1 [astro-ph.SR], arxiv.org/abs/2201.07770

© 2022 Science X Network

Citation: Study inspects the origin of high-mass X-ray binary 4U 2206+54 (2022, January 31)

retrieved 2 May 2024 from <https://phys.org/news/2022-01-high-mass-x-ray-binary-4u.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.