

A link between early maturation and better aerobic performance identified in juvenile salmon

January 28 2022



Juvenile salmon. Credit: Mikko Kytökorpi

A study conducted at the University of Helsinki indicates that early sexual maturation and high aerobic performance in salmon have a

genetic link that is already evident in juvenile salmon.

Salmon are born in rivers, migrate to the sea to reach maturity and return to spawn in their native river. While the [salmon](#) that reach their maturity at an early age return from the sea after a single year, the individuals that are slower to develop can spend two years or an even longer time at sea.

Genomic regions have been identified in the salmon genome that heavily regulate the number of years spent at sea. Under the direction of Academy of Finland Research Fellow Tutku Aykanat, researchers from the University of Helsinki bred a large number of young salmon that differed on the basis of two such regions.

The researchers measured the basal metabolism and aerobic performance of the [juvenile salmon](#) before the onset of their [maturation](#) process. Carriers of gene variants associated with early [sexual maturation](#) were found to have better aerobic performance than those whose variants were associated with the late onset of sexual maturity. Aerobic performance denotes the metabolic capacity to produce energy through aerobic cellular respiration, for example, in muscles.

"Greater aerobic performance can promote early maturation, since growth, [food supply](#) and reproduction require energy produced through aerobic metabolism," says Postdoctoral Researcher Jenni Prokkola from the Faculty of Biological and Environmental Sciences, University of Helsinki.

The genetic coupling of age-at-maturity and performance did not depend on the amount of food available to the salmon, which indicates that the finding could be generalisable to both wild salmon populations and fish farming conditions.

"Salmon that have spent several years at sea and mature at a later date

are considerably larger and produce a larger number of offspring when they return to spawn compared to salmon that spawn after only one year at sea. Now, it would be important to determine whether these salmon are more susceptible to [global warming](#) due to their poorer aerobic performance. Higher water temperatures increase the energy needs of fish, but the limitations of aerobic performance make adaptation to high temperatures challenging. Aerobic performance can become an increasingly important factor for salmon in warming habitats," Prokkola says.

Investigating the genetic and physiological factors that affect age at sexual maturity is important for understanding changes taking place in salmon populations.

The research was published in *Proceedings of the Royal Society B: Biological Sciences*.

More information: Jenni M. Prokkola et al, Genetic coupling of life-history and aerobic performance in Atlantic salmon, *Proceedings of the Royal Society B: Biological Sciences* (2022). [DOI: 10.1098/rspb.2021.2500](#)

Provided by University of Helsinki

Citation: A link between early maturation and better aerobic performance identified in juvenile salmon (2022, January 28) retrieved 3 May 2024 from <https://phys.org/news/2022-01-link-early-maturation-aerobic-juvenile.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.