

Researchers develop new method to increase effectiveness of nanomedicines

January 11 2022



Credit: CC0 Public Domain

Researchers at Penn Medicine have discovered a new, more effective method of preventing the body's own proteins from treating nanomedicines like foreign invaders, by covering the nanoparticles with



a coating to suppress the immune response that dampens the therapy's effectiveness.

When injected into the bloodstream, unmodified nanoparticles are swarmed by elements of the immune system called <u>complement proteins</u>, triggering an <u>inflammatory response</u> and preventing the nanoparticles from reaching their therapeutic targets in the body. Researchers have devised some methods to reduce this problem, but the Penn Medicine team, whose findings are published in *Advanced Materials*, has invented what may be the best method yet: coating nanoparticles with natural suppressors of complement activation.

Nanoparticles are tiny capsules, typically engineered from proteins or fatrelated molecules, that serve as delivery vehicles for certain types of treatment or vaccine—usually those containing RNA or DNA. The bestknown examples of nanoparticle-delivered medicines are mRNA vaccines against COVID-19.

"It turned out to be one of those technologies that just works right away and better than anticipated," said study co-senior author Jacob Brenner, MD, Ph.D., an associate professor of Pulmonary Medicine in the Division of Pulmonary, Allergy, and Critical Care.

The Complement Problem

Therapies based on RNA or DNA generally need delivery systems to get them through the bloodstream into target organs. Harmless viruses often have been used as carriers or "vectors" of these therapies, but nanoparticles are increasingly considered safer alternatives. Nanoparticles also can be tagged with antibodies or other molecules that make them hone in precisely on targeted tissues.

Despite its promise, nanoparticle-based medicine has been greatly



limited by the complement attack problem. Circulating complement proteins treat nanoparticles as if they were bacteria, immediately coating nanoparticle surfaces and summoning large white blood cells to gobble up the "invaders." Researchers have attempted to reduce the problem by pre-coating nanoparticles with camouflaging molecules—for example, the organic compound polyethylene glycol (PEG) attracts water molecules to form a watery, protective shell around nanoparticles. But nanoparticles camouflaged with PEG or other protective substances still draw at least some complement attack. In general, nanoparticle-based medicines that must move through the bloodstream to do their work (mRNA COVID-19 vaccines are injected into muscle, not the bloodstream) have had a very low efficiency in getting to their target organs, usually less than one percent.

Borrowing a Strategy

In the study, Brenner and Myerson and their team came up with an alternative or add-on approach to protect nanoparticles—an approach based on natural complement-inhibitor proteins that circulate in the blood, attaching to human cells to help protect them from complement attack.

The researchers found that, in lab-dish experiments, coating standard PEG-protected nanoparticles with one of these complement inhibitors, called Factor I, provided dramatically better protection from complement attack. In mice, the same strategy prolonged the half-life of standard nanoparticles in the bloodstream, allowing a much larger fraction of them to reach their targets.

"Many bacteria also coat themselves with these factors to protect against complement attack, so we decided to borrow that strategy for nanoparticles," said co-senior author Jacob Myerson, Ph.D., a senior research scientist in the Department of Systems Pharmacology and



Translational Therapeutics at Penn.

In a set of experiments in mouse models of severe inflammatory illness, the researchers also showed that attaching Factor I to nanoparticles prevents the hyper-allergic reaction that otherwise could be fatal.

Further testing will be needed before nanomedicines incorporating Factor I can be used in people, but in principle, the researchers said, attaching the complement-suppressing <u>protein</u> could make nanoparticles safer and more efficient as therapeutic delivery vehicles so that they could be used even in severely ill patients.

The researchers now plan to develop strategies for protecting not only nanomedicines but also medical devices, such as catheters, stents and dialysis tubing, which are similarly susceptible to complement attack. They also plan to investigate other protective proteins beside Factor I.

"We're recognizing now that there's a whole world of proteins that we can put on the surface of <u>nanoparticles</u> to defend them from immune attack," Brenner said.

More information: Zhicheng Wang et al, Combating Complement's Deleterious Effects on Nanomedicine by Conjugating Complement Regulatory Proteins to Nanoparticles, *Advanced Materials* (2021). DOI: 10.1002/adma.202107070

Provided by Perelman School of Medicine at the University of Pennsylvania

Citation: Researchers develop new method to increase effectiveness of nanomedicines (2022, January 11) retrieved 8 May 2024 from https://phys.org/news/2022-01-method-effectiveness-



nanomedicines.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.