

# Researchers reveal preservation mechanism of *Chuar* fossils in Lantian biota

January 7 2022, by Li Yuan

---



China Anhui Wannan, large tracts of agricultural land. Credit: Pixabay

A research group led by Dr. Wang Wei from the Nanjing Institute of Geology and Palaeontology of the Chinese Academy of Sciences (NIGPAS) employed light microscopy, field emission scanning electron

microscopy, energy-dispersive spectroscopy, and Raman spectroscopy techniques to reveal the preservation mechanism of pyritized *Chuarina* from the Lantian biota.

The study was published in *Precambrian Research* on Dec. 31.

The Lantian biota (approximately 600 Ma) was discovered from the Ediacaran basinal-facies deposition of the Lantian Formation near Lantian Town of the Anhui Province, South China. The *Chuarina* fossils, usually with spheroidal or disk-like shapes, are the most common macrofossils in the Lantian biota.

Although great majority of the Lantian *Chuarina* fossils are preserved as carbonaceous compressions, *Chuarina* has been reported to be commonly preserved as pyritization enveloped by aluminosilicate minerals in the upper Member II of the Lantian Formation.

The outer minerals that envelope the pyrite layer have been assigned to quartz or aluminosilicate [clay minerals](#). However, their exact mineral compositions as well as whether these minerals have contributed to the exceptional preservation of Ediacaran Lantian *Chuarina* remain unclear.

"Our taphonomic experiments show that *Chuarina* fossil pyritization occurred at early fossilization process and had contributed to the soft-tissue preservation," said Dr. Wang. "The pyritization process was probably divided into at least two stages, which were characterized by distinct pyrite crystal morphologies."

The pyritized *Chuarina* fossils were enveloped by platy minerals, a complex mixture of quartz and magnesium-rich aluminosilicates. Integrated [mineral](#) and structural pattern analyses showed that these quartz and clay minerals were both secondary overgrowth on *Chuarina* internal mold, which probably formed as a diagenetic product concurrent

with carbonate dissolution in the Lantian black shales.

Their formation was probably regulated by local micro-environment near the Chuaria bodies. These enveloped minerals might facilitate preservation and identification of Chuaria fossils, but they were probably not involved in the initial fossilization process as the early diagenetic pyritization had done.

**More information:** Wei Wang et al, Taphonomic study of Chuaria fossils from the Ediacaran Lantian biota of South China, *Precambrian Research* (2021). [DOI: 10.1016/j.precamres.2021.106529](https://doi.org/10.1016/j.precamres.2021.106529)

Provided by Chinese Academy of Sciences

Citation: Researchers reveal preservation mechanism of Chuaria fossils in Lantian biota (2022, January 7) retrieved 28 June 2024 from <https://phys.org/news/2022-01-reveal-mechanism-chuarial-fossils-lantian.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.