

Plants: RNA notes to self

January 13 2022



An image of an *Arabidopsis thaliana* plant, a relative of mustard, developing properly. Cold Spring Harbor Laboratory Professor David Jackson and his team discovered a protein required for normal development that helps plant cells exchange RNA messages. Credit: Munenori Kitagawa/Jackson lab/CSHL, 2021

How does a [developing plant shoot](#) know how, where, and when to grow? Dividing cells need to pass messages from one another to coordinate growth. In plants, important messages are packaged into RNA, which are sent from cell to cell. By studying the mustard-like

plant *Arabidopsis thaliana*, Cold Spring Harbor Laboratory (CSHL) Professor David Jackson and his team found that RNA messages need a special protein to escort them where they need to go. Without this escort, cells cannot coordinate and the plant fails to develop properly.

Unlike [animal cells](#), plant [cells](#) are surrounded by a rigid cell wall. Messages can cross this wall through tiny holes called plasmodesmata. Munenori Kitagawa, a postdoc in the Jackson lab who led this study, says, "Plasmodesmata are nanochannels embedded in the cell wall. They mediate various signals' transport from cell to cell, including [protein](#), RNA, hormones, ions, and nutrients."

Kitagawa wondered how the plasmodesmata's gates regulate messaging from one cell to the next. The team discovered that RNA signaling relied on a protein called AtRRP44a. Lowering the amount of AtRRP44a slowed the movement of RNA messages; lacking the right messages, the plants failed to develop properly. A protein similar to this escort protein is present in other plants, yeast, and animals. The researchers were able to swap out part of the *Arabidopsis thaliana* signaling system with parts from corn and restore [normal development](#), showing that this signaling system is similar in many kinds of plants. Jackson says, "Plants are very sophisticated. We think of them just sitting in their environment, not moving, but really they're processing a lot of information. The different parts of the plant are talking to each other, sharing whether they have some pathogen attack or if they need some nutrients."



Defects in an RNA-based cell-to-cell RNA messaging system result in poor development, such as in this *Arabidopsis thaliana* plant. Credit: Kitagawa/Jackson lab/CSHL, 2021

In a related [study published recently in the journal *Science*](#), Jackson and collaborators at New York University found that signals transported

through these gates can increase the number of cell layers in corn roots, making the [plants](#) potentially more resilient to environmental changes.

"This paper represents an important step towards understanding how information is exchanged between cells to control development and other processes," said John McDowell, a program officer in the U.S. National Science Foundation's Directorate for Biological Sciences. "By revealing a new component of cell-to-cell communication, this research opens the door for further investigation that could allow us to harness this process."

More information: Munenori Kitagawa et al, An RNA exosome subunit mediates cell-to-cell trafficking of a homeobox mRNA via plasmodesmata, *Science* (2022). [DOI: 10.1126/science.abm0840](https://doi.org/10.1126/science.abm0840).
www.science.org/doi/10.1126/science.abm0840

Provided by Cold Spring Harbor Laboratory

Citation: Plants: RNA notes to self (2022, January 13) retrieved 2 May 2024 from <https://phys.org/news/2022-01-rna.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--