

## Scientists make a new type of optical device using alumina

January 27 2022



Figure 1: A schematic diagram showing how researchers can study radiation from the early universe using an infrared reflective filter with a moth eye structure created by a pulsed laser process. Credit: NASA, ESA, M. Postman (STScI), and the CLASH Team / NRAO, AUI, NSF / modified by Kavli IPMU

## Scientists from the Kavli Institute for the Physics and Mathematics of



the Universe and the University of Minnesota, Tomotake Matsumura and Shaul Hanany, and their collaborators have made a new type of optical element that will improve the performance of telescopes studying radiation from the Big Bang.

The <u>cosmic microwave background</u> (CMB) is a relic radiation remnant from the big bang. It reaches our telescopes after traveling 14 billion years since the birth of the Universe. Studying the properties of this radiation, scientists infer the physics of the <u>big bang</u>, how clusters of galaxies form, and the matter and energy content in the Universe. Four Nobel prizes have been awarded for past studies of the CMB.

To study the CMB, telescopes must be tuned to wavelengths in which it is most intense, about 1-3 mm, and they must separate out shorter wavelength radiation that the atmosphere and Milky Way emit. Among the most effective optical elements that absorbs the short wavelength radiation but lets the CMB pass through is alumina, a material made of aluminum and oxygen and that is second in hardness only to diamond. One challenge with using alumina is that it also reflects almost 50% of the radiation impinging on it. Matsumura and Hanany have now come up with a new way to fabricate anti-reflective structures that reduce reflections fifty-fold.





Figure 2: Images of one side of the fabricated filter. The other side is identical. (a) A photograph of the entire filter. The ruler is graduated up to a length of 300 mm. (b) An enlarged area. (c) Rendering of confocal microscopy scanning of the SWS. Credit: Takaku et al.

The researchers partnered with Mark Devlin and Simon Dicker, colleagues at the University of Pennsylvania who are operating the MUSTANG2 instrument, which is coupled to the Green Bank Telescope in Virginia. Hanany and Matsumura provided the MUSTANG2 team with an alumina short-wavelength absorber that has the new antireflective structures. The MUSTANG2 instrument is now conducting sky observations with the new technology, demonstrating for the first time its success.





Figure 3: The transmission spectrum for the MUSTANG2 filter for unpolarized light (upper panel, blue dots), for s- and p-polarized light (lower panel, red and green dots, respectively), and RCWA predicted transmission given the average shape data for each side as provided in Table 2 (upper panel, solid cyan, and lower panel solid red and green). The average transmission within the band (white region) is 98%. Note the different vertical spans for the two panels, which provide both an overall performance view and details on the agreement between the data and the RCWA predictions. Credit: Takaku et al.



The researchers patterned the alumina with small pyramidal structures, which are about one millimeter tall (0.04 inch) and repeat across the 30 cm (one foot) diameter with a periodicity of just less than one millimeter. It has long been known that incorporating such structures on the surfaces of materials reduces reflections. With the small pyramids, light enters and is leaving the material more gradually, leading to much lower reflection. Matsumura and Hanany's innovation is in the way they patterned the alumina, which is too hard to be machined with standard tools. They used ultra-short pulsed laser, with pulses few trillionths of a second long and reaching 100 megawatts each, to ablate the material away and to shape the surface relief to its optimal anti-reflective shape. Within about four days the laser process produced 320,000 pyramids on both sides of the alumina disc. The researchers measured the properties of the alumina sample and showed that it reflects less than 1% of the incident <u>radiation</u>. This is the first time such an optical element has been fabricated and coupled to an operating instrument, and it is the largest sample of <u>alumina</u> to have been laser-ablated.

This innovation will lead to more efficient instruments peering back in time and revealing the physical process at the Big Bang and throughout the evolution of the universe.

**More information:** Ryota Takaku et al, A Large Diameter Millimeter-Wave Low-Pass Filter Made of Alumina with Laser Ablated Anti-Reflection Coating, *Optics Express* (2021). <u>DOI: 10.1364/OE.444848</u>

Provided by Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU)



Citation: Scientists make a new type of optical device using alumina (2022, January 27) retrieved 1 May 2024 from <u>https://phys.org/news/2022-01-scientists-optical-device-alumina.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.