

'Slushy' magma ocean led to formation of the Moon's crust

January 13 2022



Side view of the crater Moltke taken from Apollo 10. Credit: Public Domain

Scientists have shown how the freezing of a 'slushy' ocean of magma may be responsible for the composition of the Moon's crust.

The scientists, from the University of Cambridge and the Ecole normale supérieure de Lyon, have proposed a new model of crystallization, where crystals remained suspended in liquid magma over hundreds of millions of years as the lunar 'slush' froze and solidified. The results are reported in the journal *Geophysical Review Letters*.

Over fifty years ago, Apollo 11 astronauts collected samples from the lunar Highlands. These large, pale regions of the Moon—visible to the naked eye—are made up of relatively light rocks called anorthosites. Anorthosites formed early in the history of the Moon, between 4.3 and 4.5 billion years ago.

Similar anorthosites, formed through the crystallization of magma, can be found in fossilized magma chambers on Earth. Producing the large volumes of anorthosite found on the Moon however, would have required a huge global magma [ocean](#).

Scientists believe that the Moon formed when two protoplanets, or embryonic worlds, collided. The larger of these two protoplanets became the Earth, and the smaller became the Moon. One of the outcomes of this collision was that the Moon was very hot—so hot that its entire mantle was molten magma, or a magma ocean.

"Since the Apollo era, it has been thought that the lunar crust was formed by light anorthite crystals floating at the surface of the liquid

magma ocean, with heavier crystals solidifying at the [ocean floor](#)," said co-author Chloé Michaut from Ecole normale supérieure de Lyon. "This 'flotation' model explains how the lunar Highlands may have formed."

However, since the Apollo missions many lunar meteorites have been analyzed and the surface of the Moon has been extensively studied. Lunar anorthosites appear more heterogenous in their composition than the original Apollo samples, which contradicts a flotation scenario where the liquid ocean is the common source of all anorthosites.

The range of anorthosite ages—over 200 million years—is difficult to reconcile with an ocean of essentially [liquid magma](#) whose characteristic solidification time is close to 100 million years.

"Given the range of ages and compositions of the anorthosites on the Moon, and what we know about how crystals settle in solidifying magma, the lunar crust must have formed through some other mechanism," said co-author Professor Jerome Neufeld from Cambridge's Department of Applied Mathematics and Theoretical Physics.

Michaut and Neufeld developed a mathematical model to identify this mechanism.

In the low lunar gravity, the settling of crystal is difficult, particularly when strongly stirred by the convecting magma ocean. If the crystals remain suspended as a crystal slurry, then when the crystal content of the slurry exceeds a critical threshold, the slurry becomes thick and sticky, and the deformation slow.

This increase of crystal content occurs most dramatically near the surface, where the slushy magma ocean is cooled, resulting in a hot, well-mixed slushy interior and a slow-moving, crystal rich lunar 'lid'.

"We believe it's in this stagnant 'lid' that the lunar crust formed, as lightweight, anorthite-enriched melt percolated up from the convecting crystalline slurry below," said Neufeld. "We suggest that cooling of the early magma ocean drove such vigorous convection that crystals remained suspended as a slurry, much like the [crystals](#) in a slushy machine."

Enriched lunar surface rocks likely formed in magma chambers within the lid, which explains their diversity. The results suggest that the timescale of [lunar crust](#) formation is several hundreds of million years, which corresponds to the observed ages of the lunar anorthosites.

Serial magmatism was initially proposed as a possible mechanism for the formation of lunar anorthosites, but the slushy model ultimately reconciles this idea with that of a global lunar [magma](#) ocean.

More information: Formation of the lunar primary crust from a long-lived slushy magma ocean, *Geophysical Review Letters*, [agupubs.onlinelibrary.wiley.co ... 10.1029/2021GL095408](https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021GL095408)

Provided by University of Cambridge

Citation: 'Slushy' magma ocean led to formation of the Moon's crust (2022, January 13) retrieved 6 June 2024 from <https://phys.org/news/2022-01-slushy-magma-ocean-formation-moon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.