

Southern Ocean storms cause outgassing of carbon dioxide

January 25 2022



A wave glider was one of the ocean robots in the project, collecting data from the surface. Credit: Fred Fourie



Storms over the waters around Antarctica drive an outgassing of carbon dioxide into the atmosphere, according to a new international study with researchers from the University of Gothenburg. The research group used advanced ocean robots for the study, which provides a better understanding of climate change and can lead to better global climate models.

The world's southernmost <u>ocean</u>, the Southern Ocean that surrounds Antarctica, plays an important role in the <u>global climate</u> because its waters contain large amounts of <u>carbon dioxide</u>. A new international study, in which researchers from the University of Gothenburg participated, has examined the complex processes driving air-sea fluxes of gasses, such as carbon dioxide.

Storms bring carbon dioxide-rich waters to the surface

The research group is now delivering new findings that shed light on the area's important role in climate change.

"We show how the intense storms that often occur in the region increase ocean mixing and bring carbon dioxide-rich waters from the deep to the surface. This drives an outgassing of carbon dioxide from the ocean to the atmosphere. There has been a lack of knowledge about these complex processes, so the study is an important key to understanding the Southern Ocean's significance for the climate and the global carbon budget", says Sebastiaan Swart, professor of oceanography at the University of Gothenburg and co-author of the study.





One of the ocean robots in the project, a slocum glider, was able to to collect data at a depth of one kilometer. Credit: Sea Technology Services

Facilitates better climate models

Half of all carbon dioxide bound in the world's oceans is found in the Southern Ocean. At the same time, climate change is expected to result in more intense storms in the future. Therefore, it is vital to understand the storms' impact on the outgassing of <u>carbon</u> dioxide into the atmosphere, the researchers point out.

"This knowledge is necessary to be able to make more accurate



predictions about future climate change. Currently, these environmental processes are not captured by global climate models", says Marcel du Plessis at the University of Gothenburg, who also participated in the study.

Pioneering ocean robotics

Measuring the inaccessible and stormy waters around Antarctica for a long period of time is a real challenge, which the researchers tackled with the help of unique robot technology. For several months, autonomous ocean robots; drones and ocean gliders, collected data from the surface and through to depths of one kilometer.

"This pioneering technology gave us the opportunity to collect data with long endurance, which would not have been possible via a research vessel. Thanks to these ocean robots we can now fill important knowledge gaps and gain a better understanding of the importance of the ocean for the <u>climate</u>, says Sebastiaan Swart.

The research was published in Nature Communications.

More information: Nicholson, SA. et al, Storms drive outgassing of CO2 in the subpolar Southern Ocean, *Nat Commun* (2022). doi.org/10.1038/s41467-021-27780-w

Provided by University of Gothenburg

Citation: Southern Ocean storms cause outgassing of carbon dioxide (2022, January 25) retrieved 26 April 2024 from https://phys.org/news/2022-01-southern-ocean-storms-outgassing-carbon.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.