

Measuring carbon ion loss from the Martian atmosphere

February 8 2022, by Morgan Rehnberg



A computer simulation shows ions escaping the Martian atmosphere due to interactions with the Sun's solar wind. Credit: <u>NASA's Scientific Visualization</u> <u>Studio</u>

In recent decades, planetary scientists have developed ample evidence indicating that Mars was home to a relatively thick atmosphere and hosted liquid water on its surface during the Noachian, around 4 billion years ago. Since then, significant climate change occurred, resulting in



the loss of much of the planet's atmosphere and all of its surface water.

What remains of the <u>atmosphere</u> today is a thin envelope consisting largely of <u>carbon dioxide</u>. Without a protective planetary magnetic field, this, too, is being stripped away by the solar wind. Since its arrival in 2014, NASA's Mars Atmosphere and Volatile Evolution (MAVEN) spacecraft has tracked the escape of the atmosphere's most abundant ions, those of atomic and <u>molecular oxygen</u>.

In a new study, Pickett et al. attempt to extend these observations to <u>carbon</u>, the second most abundant element in the atmosphere. A complicating factor in observing the atmosphere's carbon (C) ion loss is that carbon is similar in mass to the dominant species, oxygen (O). For this work, the authors use 40 orbits of data collected by MAVEN's Suprathermal and Thermal Ion Composition (STATIC) instrument, which records the charge, mass, and velocity of incoming ions.

Because of instrumental uncertainties, STATIC registers particle impacts with a small range of masses; the distribution is centered on the true mass of a given ion. Because C+ and O+ have similar masses, these distributions overlap. Thus, contributions from the oxygen ions must be modeled and removed before the C+ population can be ascertained. The authors develop and describe a multistep process to accomplish this.

The researchers applied their method to data collected by STATIC in April 2018 as an initial assessment of the technique and a first approximation of the escape rate of C+. They estimate that C+ flows away from Mars with a flux on the order of 104–105 per centimeter squared per second down the magnetotail. This finding is broadly consistent with previous model-derived estimates of carbon loss from the atmosphere and lower by 1–2 orders of magnitude than the observed loss rate for oxygen ions.



In aggregate, STATIC found detectable C+ fluxes in about a quarter of all observations during the study period. This illustrates the persistent difficulty of disentangling the two most abundant components of the Martian atmosphere. Understanding the related, yet separate, stories of carbon and <u>oxygen</u> in the current atmosphere will be a future task with implications for the history of the red planet's climate.

More information: N. B. Pickett et al, Carbon Ion Fluxes at Mars: First Results of Tailward Flows From MAVEN-STATIC, *Journal of Geophysical Research: Space Physics* (2021). DOI: 10.1029/2021JA029635

This story is republished courtesy of Eos, hosted by the American Geophysical Union. Read the original story <u>here</u>.

Provided by American Geophysical Union

Citation: Measuring carbon ion loss from the Martian atmosphere (2022, February 8) retrieved 26 June 2024 from <u>https://phys.org/news/2022-02-carbon-ion-loss-martian-atmosphere.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.