

# **Experimental evidence for long-distance electrodynamic intermolecular forces**

February 28 2022, by Thamarasee Jeewandara





Long-range ED interactions: Principle and experimental approaches. (A) At thermal equilibrium, macromolecules show a Brownian diffusive motion in solution (left). By switching-on an external energy source, molecules are in an out-of-thermal equilibrium collective vibrational state that can generate ED forces through associated large dipolar resonant oscillations (right). (B) In THz spectroscopy, the frequency of the collective vibration varies as a function of the average intermolecular distance (determined by the protein concentration). With respect to the reference frequency at "infinite" dilution, a frequency shift inversely proportional to the cubic power of the average intermolecular distance is theoretically expected if the proteins interact through ED forces. (C) In FCS, the transit times of the proteins across the volume of observation are measured and, hence, the protein diffusion coefficients. Under out-of-thermal equilibrium conditions, the theory predicts a phase transition due to ED forces clustering proteins. This should be observed at a given protein concentration by a sudden drop in the diffusion coefficient from its Brownian value. In both (B) and (C), out-of-thermal equilibrium is activated by optical excitation. Credit: Science Advances, DOI: 10.1126/sciadv.abl5855

While classical and quantum <u>electrodynamics</u> present the existence of dipole-dipole long-range electrodynamics forces, they remain to be experimentally observed. The discovery of completely new and unanticipated forces that are present between biomolecules have considerable impact to understand the dynamics of molecular machines at work within living organisms. In a new report now published in *Science Advances*, Mathias Lechelon and a research team at the French National Centre for Scientific Research (CNRS) France conducted two independent experiments based on different physical effects, which they detected via <u>fluorescence correlation spectroscopy</u> and <u>terahertz</u> <u>spectroscopy</u>, respectively, to demonstrate experimental activation of resonant electrodynamic intermolecular forces. The outcomes provided unprecedented experimental proof-of-principle of a physical phenomenon with importance in biology. According to the study, aside from thermal fluctuations that randomly drove molecular motion,



resonant and selective electrodynamic forces contributed to molecular encounters in crowded cellular spaces.

## Electrodynamic (ED) forces

Physicists are interested in observing intermolecular dipole-dipole electrodynamic (ED) forces, and in this study, Lechelon et al. detected these forces at the molecular level. From a physicist's viewpoint, living matter offers a range of fascinating dynamic phenomena, including biomolecules arranged in intricate and complex networks for biochemical interactions to provide astonishing efficiency. Numerically, human cells contain about 130,000 binary interactions between proteins, and about 34,000 unique human proteinprotein interactions are listed on databases. One of the most notable challenges in understanding biochemistry from a physics standpoint is to comprehend how the right molecule gets to the accurate place, at the exact moment in time, within the correct cascade of biochemical events during any biological action. Based on existing fundamental interactions, dipole-dipole interactions are predicted to promote molecular attraction due to selective resonance across long distances. In this work, Lechelon et al. experimentally determined the activation of long-range electrodynamic forces between proteins.





Collective oscillation of the R-PE measured by THz spectroscopy. Absorption strength of the R-PE (1 mg/ml in 200 mM NaCl) as a function of detection frequency and the laser power activating the molecular collective oscillation at a frequency of 71 GHz (2.4 cm-1). The absorption line profile (bottom) indicated by the horizontal white dashed line was measured at an optical power of 31.5 mW, while the laser power threshold (left) behavior is indicated by the black dashed line. Stars stand for experimental data, and the interpolating line is just a guide to the eye. Credit: *Science Advances*, DOI: 10.1126/sciadv.abl5855

#### Measuring electrodynamic forces in living matter: The experiments



To experimentally determine the activation of long-range electrodynamics forces between proteins, they used <u>R-phycoerythrin</u>; a <u>protein</u> that can be naturally excited via an external energy supply such as a light source. They then worked at different concentrations including intermolecular distances, under the excitation power of a laser <u>light</u> source to observe the activated collective intramolecular oscillation of proteins and their physical mechanisms. During the study, Lechelon et al. used two experimental methods, <u>terahertz spectroscopy</u> with a terahertz rectenna or a microwire probe as sensors and <u>fluorescence</u> <u>correlation spectroscopy</u>. The experimental work supported a proof-ofprinciple on the capacity for collective oscillations to activate dipoledipole electrodynamic intermolecular forces to explore the potential of electrodynamic forces in living matter.





R-phycoerythrin (R-PE) coherent vibrational states. Comparison of the R-PE in saline solution (red) and saline solution without R-PE (black). Two collective extension modes of R-PE appear at 71 and 96 GHz. Experimental data are indicated by full circles, and Lorentz fit is indicated by a solid line. AU, arbitrary units. Credit: *Science Advances*, DOI: 10.1126/sciadv.abl5855

# **Analyzing R-phycoerythrin**

During the initial experiments, Lechelon et al. used a natural lightharvesting protein derived from red algae, known as R-phycoerythrin.



The hexamer protein contained an  $\alpha$  subunit of two phycoerythrobilins, a  $\beta$  subunit containing three phycoerythrobilins and one phycourobilin and a  $\gamma$  subunit with two or three phycoerythrobilins or one or two phycourobilins. All these subunits contributed to the formation of one of the brightest fluorescent dyes with a sensitivity of 488 nm wavelength of light. Under stable illumination, the molecules entered a coherent vibrational state in saline solution. In the saturation regime, the team noted two collective oscillation frequencies for the protein: one at 0.071 THZ and the other at 0.096 THZ. The team schematized R-phycoerythrin as a torus-shaped object with appropriate parameters computed, including its density and Young's modulus.

# **Observing phase transitions**

The scientists next obtained a theoretically expected signature of attractive electrodynamic forces among macromolecules vibrating at the same frequency in order to show a phase transition between rapidly diffusing molecules and a clustered phase of very slow moving aggregates of molecules. The team regulated this transition via the average intermolecular distance by adjusting the molecular concentration. During <u>fluorescence correlation spectroscopy</u> experiments, Lechelon et al. varied the protein concentrations and retained the ionic strength of the protein solution at 200 mM via suitable concentrations of salt in water. This ensured good shielding of electrostatic interactions, then by varying the laser power, they repeated typical fluorescence traces for different laser powers and protein concentrations.







Effect of protein concentration and laser power illumination on R-PE diffusion: Clustering phase transition. (A) Diffusion coefficients normalized to the Brownian D0 values measured for each data series at 0.223  $\mu$ M ( $\langle r \rangle \simeq 1950$  Å) and recorded at 50 (blue circles), 100 (green triangles), and 150  $\mu$ W (red triangles). Each point corresponds to the average of five independent experiments. (B) Molecular dynamics computation of self-diffusion coefficient D/D0 (normalized to the Brownian value D0) versus intermolecular average distance  $\langle d \rangle$  for a system of particles in a cubic box interacting through longrange ED forces. Credit: *Science Advances*, DOI: 10.1126/sciadv.abl5855

# **Diffusion coefficients**

By measuring the diffusion times from cross-correlation functions of fluorescent traces and by measuring the confocal volume where the molecules are both excited and observed, the team estimated the diffusion coefficients. They then normalized the values of diffusion coefficient relative to Brownian values at a given laser power. Thereafter, the scientists showed how the measured values of diffusion coefficient matched the expected theoretical values at low laser power. For example, at a laser power of 50  $\mu$ W, the observed diffusion coefficient did not change with the intermolecular distance. For all concentrations considered, the diffusion of R-phycoerythrin molecules were **Brownian**. Lechelon et al. further conducted molecular dynamics simulations to estimate the effect of long-range electrodynamics interactions for a system of interacting molecules to allow direct comparisons between the outcome of numerical simulations and the outcome of lab experiments. The team noted the appearance of clusters that were theoretically predicted via sudden enhancement of fluorescence fluctuation amplitudes and long transit times using fluorescence spectroscopy.









Frequency shifts of the intramolecular collective vibrations of R-PE and BSA at different concentrations. Measurements were performed at room temperature in aqueous solution with 200 mM of NaCl. (A) R-PE. The shift is relative to the reference frequency measured at the lowest protein concentration. Measurements have been performed at different powers of the laser: 31.5 (blue circles), 39.5 (green triangles), and 50 mW (red triangles). Purple squares and orange stars refer to theoretical outcomes worked out with different values of molecular dipole moments. (B) BSA at a laser power of 40 mW. Insets: The frequency shifts are measured through a Lorentz fitting of the experimental resonances. The different colors in the insets are just a visual help. Credit: *Science Advances*, DOI: 10.1126/sciadv.abl5855

## **Experimental optimization**

By increasing the laser power, Lechelon et al. showed that higher the power, the stronger the oscillation of R-phycoerythrin molecules observed, along with a larger oscillating dipole moment for stronger molecular oscillations. The team could rapidly disaggregate clusters by suddenly lowering the power of the laser light supply in a clustered phase. Under the same conditions of temperature and particle concentration, when the energy input rate was below the threshold, it activated intramolecular collective vibrations, causing the disappearance of clustering transitions, to provide clear proof of the activation of electrodynamic intermolecular attractive forces via collective vibrations of the R-phycoerythrin molecules. Using further studies, Lechelon et al. confirmed the activation of long-distance electrodynamics within <u>bovine</u> serum albumin as well.





THz spectroscopy experimental setups. (A) Rectenna for the 0.07 to 0.11 THz band. (B) Microwire-based probe for the 0.25- to 0.37-THz band. Credit: *Science Advances*, DOI: 10.1126/sciadv.abl5855

## Outlook

In this way, Mathias Lechelon, and colleagues, experimentally showed the excitation of out-of-equilibrium collective oscillations to drive molecular associations via electrodynamic molecular <u>force</u> activation. The R-phycoerythrin proteins incorporated in the work provided convincing evidence for these forces based on complementary experimental approaches. The team consider the protein to be the "Rosetta stone' to identify attractive intermolecular forces.



They activated long-range electrodynamic interactions with a second protein fluorescent dye-labelled bovine serum albumin. The outcomes confirmed the presence of electrodynamic forces in the complex environment encountered in cell biology. Due to their dynamic and reversible nature, electrodynamic forces are instrumental to explain the rationale of many cellular processes. These findings are of interest to regulate the space, time of molecular interactions in living cells and to depict cellular functions. These forces have applications in the domain of drug design to improve therapeutic efficiency and the results may also shed light to overcome adverse effects of molecular overcrowding in cells.

**More information:** Mathias Lechelon et al, Experimental evidence for long-distance electrodynamic intermolecular forces, *Science Advances* (2022). DOI: 10.1126/sciadv.abl5855

Kavitha Venkatesan et al, An empirical framework for binary interactome mapping, *Nature Methods* (2008). DOI: 10.1038/nmeth.1280

© 2022 Science X Network

Citation: Experimental evidence for long-distance electrodynamic intermolecular forces (2022, February 28) retrieved 26 June 2024 from <u>https://phys.org/news/2022-02-experimental-evidence-long-distance-electrodynamic-intermolecular.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.