

# Larvicidal flavonoids inhibit key enzyme in yellow fever mosquitoes

February 17 2022

---



Credit: Charles Parker from Pexels

When most people think of flavonoids, natural compounds found in plants and other organisms, their nutritional benefits probably come to mind first. But these compounds may have another health benefit:

Researchers from Japan have discovered that certain flavonoids inhibit development in mosquitoes that can spread disease.

In a study published this month in *BMC Biology*, researchers from the University of Tsukuba have revealed that particular flavonoids inhibit an enzyme involved in the formation of a key insect hormone in the yellow fever mosquito, *Aedes aegypti*.

Mosquito-borne diseases are a major component of the worldwide burden of infectious disease in humans. *Aedes aegypti* is from a group of [mosquitoes](#) that can spread a number of viruses that cause infectious diseases in humans, including dengue fever, yellow fever, and Zika. In the wild, *A. aegypti* has begun to show resistance to insecticides, revealing a need for new types of pesticides for targeting this species.

"Flavonoids—a type of metabolic product from plants, fungi, and other organisms—can interfere with insect development and physiology, and have the ability to kill larvae of *A. aegypti*," says senior author of the study, professor Ryusuke Niwa. "Flavonoids are thought to be relatively safe for the environment, as well as human and animal health."

To investigate how flavonoids can kill [mosquito larvae](#), the researchers analyzed the activities of several flavonoids in *A. aegypti*, including daidzein, which has previously been identified as a larvicide for this species. The team found that the flavonoids inhibit the activity of glutathione S-transferase Noppera-bo (Nobo); in *A. aegypti*, Nobo is an enzyme involved in the biosynthesis of the hormone ecdysone.

Ecdysone is an insect steroid hormone, or ecdysteroid, required for the initiation of metamorphosis and regulation of molting. Because ecdysteroids are key to the life cycle of insects, chemical inhibitors of enzymes involved in making these hormones, including Nobo, are thought to be insect growth regulators (IGRs) that disrupt development

in insects without affecting other organisms.

"We also discovered that, of the flavonoids we tested, desmethylglycitein (DMG) was the most efficient Nobo inhibitor in this species, even more so than daidzein," says professor Niwa. "DMG showed larvicidal activity against *A. aegypti*, and indicated promise for DMG-based insecticides in the future."

The high prevalence of resistance in mosquitoes to current insecticides in some areas urgently requires the development of new insecticides with different chemical structures and targeting pathways from those currently in use. The results of this study offer a new avenue for developing new IGRs that are environmentally friendly and can be used for the control of mosquito populations by inhibiting the biosynthesis of ecdysteroids.

The article, "Molecular action of larvicidal flavonoids on ecdysteroidogenic glutathione S-transferase Noppera-bo in *Aedes aegypti*," was published in *BMC Biology*.

**More information:** Kazue Inaba et al, Molecular action of larvicidal flavonoids on ecdysteroidogenic glutathione S-transferase Noppera-bo in *Aedes aegypti*, *BMC Biology* (2022). [DOI: 10.1186/s12915-022-01233-2](https://doi.org/10.1186/s12915-022-01233-2)

Provided by University of Tsukuba

Citation: Larvicidal flavonoids inhibit key enzyme in yellow fever mosquitoes (2022, February 17) retrieved 10 May 2024 from <https://phys.org/news/2022-02-larvicidal-flavonoids-inhibit-key-enzyme.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.