

Opportunities for natural enemy to fight devastating fall armyworm

February 14 2022



Telenomus remus provides good opportunities for the establishment of an augmentative biological control program, reinforcing sustainable production of major crops such as maize in affected countries. Credit: G. Goergen, IITA

A new CABI-led review has highlighted mass rearing techniques,

estimated costs of mass production and release strategies for the natural enemy *Telenomus remus* that suggests it could be effective in the fight against the devastating fall armyworm (*Spodoptera frugiperda*) in affected countries.

The study, published in *CABI Agriculture and Bioscience*, states how *T. Remus* has already been used against fall armyworm in the Americas for many years and is even effective against other species of the genus *Spodoptera*—all of which blight the crops of millions of [smallholder farmers](#) including the staple crop maize.

Fall armyworm is a significant pest of over 100 crops but favors maize. The recent CABI paper "Towards estimating the economic cost of invasive alien species to African crop and livestock production," for example, estimates that in Africa the pest causes annual yield losses of around USD \$9.4 billion.

Currently, in maize, the pest is predominantly controlled by pesticides or transgenic events. However, the use of biological control agents is considered the most sustainable and preferred method of control, providing high effectiveness. Among the various natural enemies reported for FAW, the egg parasitoid *Telenomus remus* has gained most interest.

Lead researcher Dr. Yelitza Colmenarez, Center Director at CABI's center in Brazil, said, "There is no ready-to-use package available to advise farmers in using *T. remus* against fall armyworm and related pests. Further studies are urgently needed to precisely determine optimal release rates, release times and frequencies, number of release points, the best stage and device for releases and other aspects such as how large the fields should be to achieve efficient pest control."

"However, high egg parasitism rates together with long-term evidence

from Venezuela suggest that *T. remus* has indeed high potential to successfully suppress fall armyworm and related pests.



Maize infested by fall armyworm. Credit: CABI

"If biological control of FAW and related pests with *T. remus* should become a viable option, release rates may need to be more closely to rates used in the 1990s in Venezuela, i.e. rates of about 5,000–10,000 wasps per ha and season, unless major breakthroughs with cheaper mass production of the parasitoid are achieved."

The review also suggests that releases for individual smallholder farmers

owning little land may be inefficient, but on the other hand that regional approaches could work very well.

The scientists believe that—similar to other biological control agents in field crops—the use of *T. remus* will best be done as a part of an Integrated Pest Management program, avoiding broad spectrum insecticides in release fields. Both quality control of the mass reared parasitoid and an optimized, cost-efficient release strategy is crucial to a successful [pest](#) management, they say.

Dr. Colmenarez added, "Due to the recent invasion of fall armyworm in Africa, Asia, and Australia, *T. remus* provides good opportunities for the establishment of an augmentative biological control program, reinforcing sustainable production of major crops such as maize in affected countries."

"When considering the use of [biological control agents](#), it is always necessary to include an assessment of possible risks for non-target effects, particularly for exotic species. In most of those countries where *T. remus* may be considered for release in the future, the parasitoid has been found to be present, e.g. China, India, Australia and several African countries, so cannot be considered exotic there."

"However, in regions where *T. remus* is not known from, its status should be carefully evaluated and appropriate risk assessment procedures followed. Altogether, *T. remus* releases may be having high prospects for contributing to FAW management in newly invaded areas though still challenges exist that would require further research."

More information: Yelitza Coromoto Colmenarez et al, The use of *Telenomus remus* (Nixon, 1937) (Hymenoptera: Scelionidae) in the management of *Spodoptera* spp.: potential, challenges and major benefits, *CABI Agriculture and Bioscience* (2022). [DOI:](#)

[10.1186/s43170-021-00071-6](https://phys.org/news/2022-02-opportunities-natural-enemy-devastating-fall.html)

Provided by CABI

Citation: Opportunities for natural enemy to fight devastating fall armyworm (2022, February 14) retrieved 16 June 2024 from <https://phys.org/news/2022-02-opportunities-natural-enemy-devastating-fall.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.