

## Trees may benefit from earlier start of growing season in cold humid areas

February 28 2022



Illustration of how warming impacts on wood phenology affect tree growth. Credit: Gao Shan

## A research group led by Prof. Liang Eryuan from the Institute of Tibetan



Plateau Research (ITP) of the Chinese Academy of Sciences has found that trees may benefit from an earlier start of the growing season in cold humid areas, but not in dry areas.

This paper was published in Nature Ecology & Evolution on Feb. 28.

Plants can naturally feel changes in their environment. When the days lengthen and the temperature warms up after winter chilling, <u>trees</u> sense that it is time to green up and reactivate growth.

Trees tend to sprout and resume growth earlier as climate warms, but whether and how an earlier start of the growing <u>season</u> influences tree growth remain unclear.

Carbon surplus from increased uptake by leaves may not mean that trees necessarily grow more. Thus, extensive observations of the growth of tree stems are essential for predicting changes in productivity because wood is the primary long-term carbon storage pool in forests.

In the study, the researchers used a large collection of annual tree-ring width data from the International Tree-Ring Data Bank, the world's largest public archive of tree ring data.

"Previous studies have shown that the onset of wood formation is closely related to the fulfillment of critical temperatures and photoperiod. Therefore, we analyzed how annual tree-ring width relates to thermal thresholds for growth start on a hemispheric scale," said Prof. Liang.

The onset of wood formation is the main factor that directly or indirectly triggers all subsequent phases of xylem maturation and determines the period of xylem growth. Small advances in the period of cell division can thus lead to substantial increases in xylem cell production, which eventually results in wider growth rings.



Therefore, tree growth in <u>cold climates</u> would be enhanced throughout the entire growing season by an earlier onset of cambial activity or by higher growth rates at the peak of the growing season.

Annual ring width is usually correlated with monthly or seasonal climate data. After extensive processing of daily climate data and conducting different analyses, the researchers found distinct spatial patterns of tree growth in response to an earlier growing season onset.

Areas where tree growth benefits from an advanced spring are generally located at the higher latitudes (above 60°N), central Europe, as well as in eastern and western coastal North America. These cool and humid regions are not strongly limited by water availability during the growing season.

The regions showing negative effects of advanced springs on tree growth were mainly located on the Colorado Plateau and the Tibetan Plateau, which correspond to cold and dry conditions, where forests are typically limited by a number of factors including low temperatures, drought events, and poor soil fertility.

"From these observations, we conclude that trees will grow faster under advanced springs, unless they are thirsty," said Prof. Liang.

In addition, the researchers analyzed the effects of advanced spring on tree growth and found distinct latitudinal responses. In boreal forests of northern Asia and Europe, advanced spring enhanced tree growth primarily due to the alleviation of cold stress. In the temperate forests of central Europe and the eastern coast of the U.S., as well as in forests of the Mediterranean region and along the western coast of the U.S., advanced spring also enhanced growth, but primarily due to the extension of the growing season.



In semi-arid forests of the Colorado Plateau and dry subalpine forests of the Tibetan Plateau, advanced spring did not benefit growth, as a longer growing season induces both atmospheric and soil drought in those regions, and also increases the risk of tree exposure to <u>spring</u> frost.

**More information:** Eryuan Liang, An earlier start of the thermal growing season enhances tree growth in cold humid areas but not in dry areas, *Nature Ecology & Evolution* (2022). DOI: 10.1038/s41559-022-01668-4. www.nature.com/articles/s41559-022-01668-4

Provided by Chinese Academy of Sciences

Citation: Trees may benefit from earlier start of growing season in cold humid areas (2022, February 28) retrieved 1 May 2024 from <u>https://phys.org/news/2022-02-trees-benefit-earlier-season-cold.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.